

PATENT COOPERATION TREATY

PCT

NOTIFICATION OF ELECTION

(PCT Rule 61.2)

From the INTERNATIONAL BUREAU

To:

Commissioner
US Department of Commerce
United States Patent and Trademark
Office, PCT
2011 South Clark Place Room
CP2/5C24
Arlington, VA 22202
ÉTATS-UNIS D'AMÉRIQUE

in its capacity as elected Office

Date of mailing (day/month/year) 09 April 2001 (09.04.01)	
International application No. PCT/EP00/06429	Applicant's or agent's file reference FC 862/5
International filing date (day/month/year) 07 July 2000 (07.07.00)	Priority date (day/month/year) 14 July 1999 (14.07.99)
Applicant MANTEGANI, Sergio et al	

1. The designated Office is hereby notified of its election made:

☒ in the demand filed with the International Preliminary Examining Authority on:

12 February 2001 (12.02.01)

☐ in a notice effecting later election filed with the International Bureau on:

2. The election ☒ was

☐ was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

The International Bureau of WIPO
34, chemin des Colombettes
1211 Geneva 20, Switzerland

Facsimile No.: (41-22) 740.14.35

Authorized officer

Nestor Santesso

Telephone No.: (41-22) 338.83.38

PATENT COOPERATION TREATY

PCT

From the INTERNATIONAL BUREAU

NOTIFICATION CONCERNING
SUBMISSION OR TRANSMITTAL
OF PRIORITY DOCUMENT

(PCT Administrative Instructions, Section 411)

To:

PHARMACIA & UPJOHN SpA
V.le Pasteur, 10
I-20014 Nerviano
ITALIE

Date of mailing (day/month/year) 16 October 2000 (16.10.00)	IMPORTANT NOTIFICATION
Applicant's or agent's file reference FC 862/5	
International application No. PCT/EP00/06429	International filing date (day/month/year) 07 July 2000 (07.07.00)
International publication date (day/month/year) Not yet published	Priority date (day/month/year) 14 July 1999 (14.07.99)
Applicant PHARMACIA & UPJOHN SpA et al	

1. The applicant is hereby notified of the date of receipt (except where the letters "NR" appear in the right-hand column) by the International Bureau of the priority document(s) relating to the earlier application(s) indicated below. Unless otherwise indicated by an asterisk appearing next to a date of receipt, or by the letters "NR", in the right-hand column, the priority document concerned was submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b).
2. This updates and replaces any previously issued notification concerning submission or transmittal of priority documents.
3. An asterisk(*) appearing next to a date of receipt, in the right-hand column, denotes a priority document submitted or transmitted to the International Bureau but not in compliance with Rule 17.1(a) or (b). In such a case, **the attention of the applicant is directed** to Rule 17.1(c) which provides that no designated Office may disregard the priority claim concerned before giving the applicant an opportunity, upon entry into the national phase, to furnish the priority document within a time limit which is reasonable under the circumstances.
4. The letters "NR" appearing in the right-hand column denote a priority document which was not received by the International Bureau or which the applicant did not request the receiving Office to prepare and transmit to the International Bureau, as provided by Rule 17.1(a) or (b), respectively. In such a case, **the attention of the applicant is directed** to Rule 17.1(c) which provides that no designated Office may disregard the priority claim concerned before giving the applicant an opportunity, upon entry into the national phase, to furnish the priority document within a time limit which is reasonable under the circumstances.

<u>Priority date</u>	<u>Priority application No.</u>	<u>Country or regional Office or PCT receiving Office</u>	<u>Date of receipt of priority document</u>
14 July 1999 (14.07.99)	9916562.3	GB	13 Sept 2000 (13.09.00)

The International Bureau of WIPO
34, chemin des Colombettes
1211 Geneva 20, Switzerland

Facsimile No. (41-22) 740.14.35

Authorized officer

Athina Nickitas-Etienne

Telephone No. (41-22) 338.83.38

PATENT COOPERATION TREATY

- 2 FEB 2001
H2

PCT

From the INTERNATIONAL BUREAU

To:

PHARMACIA & UPJOHN SpA
V.le Pasteur, 10
I-20014 Nerviano
ITALIENOTICE INFORMING THE APPLICANT OF THE
COMMUNICATION OF THE INTERNATIONAL
APPLICATION TO THE DESIGNATED OFFICES

(PCT Rule 47.1(c), first sentence)

Date of mailing (day/month/year) 25 January 2001 (25.01.01)		
Applicant's or agent's file reference FC 862/5		IMPORTANT NOTICE
International application No. PCT/EP00/06429	International filing date (day/month/year) 07 July 2000 (07.07.00)	
Priority date (day/month/year) 14 July 1999 (14.07.99)		
Applicant PHARMACIA & UPJOHN SpA et al		

1. Notice is hereby given that the International Bureau has communicated, as provided in Article 20, the international application to the following designated Offices on the date indicated above as the date of mailing of this Notice:

AU, KP, KR, US

In accordance with Rule 47.1(c), third sentence, those Offices will accept the present Notice as conclusive evidence that the communication of the international application has duly taken place on the date of mailing indicated above and no copy of the international application is required to be furnished by the applicant to the designated Office(s).

2. The following designated Offices have waived the requirement for such a communication at this time:

AE, AL, AM, AP, AT, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EA, EE, EP, ES, FI, GB, GD,
GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
NO, NZ, OA, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW

The communication will be made to those Offices only upon their request. Furthermore, those Offices do not require the applicant to furnish a copy of the international application (Rule 49.1(a-bis)).

3. Enclosed with this Notice is a copy of the international application as published by the International Bureau on
25 January 2001 (25.01.01) under No. WO 01/05756

REMINDER REGARDING CHAPTER II (Article 31(2)(a) and Rule 54.2)

If the applicant wishes to postpone entry into the national phase until 30 months (or later in some Offices) from the priority date, a demand for international preliminary examination must be filed with the competent International Preliminary Examining Authority before the expiration of 19 months from the priority date.

It is the applicant's sole responsibility to monitor the 19-month time limit.

Note that only an applicant who is a national or resident of a PCT Contracting State which is bound by Chapter II has the right to file a demand for international preliminary examination.

REMINDER REGARDING ENTRY INTO THE NATIONAL PHASE (Article 22 or 39(1))

If the applicant wishes to proceed with the international application in the national phase, he must, within 20 months or 30 months, or later in some Offices, perform the acts referred to therein before each designated or elected Office.

For further important information on the time limits and acts to be performed for entering the national phase, see the Annex to Form PCT/IB/301 (Notification of Receipt of Record Copy) and Volume II of the PCT Applicant's Guide.

The International Bureau of WIPO
34, chemin des Colombettes
1211 Geneva 20, Switzerland

Facsimile No. (41-22) 740.14.35

Authorized officer

J. Zahra

Telephone No. (41-22) 338.83.38

PATENT COOPERATION TREATY

PCT

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

Applicant's or agent's file reference FC 862	FOR FURTHER ACTION See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)	
International application No. PCT/EP00/06429	International filing date (<i>day/month/year</i>) 07/07/2000	Priority date (<i>day/month/year</i>) 14/07/1999
International Patent Classification (IPC) or national classification and IPC C07C317/46		
Applicant PHARMACIA & UPJOHN SpA et al.		


1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.
2. This REPORT consists of a total of 5 sheets, including this cover sheet.

☐ This report is also accompanied by ANNEXES, i.e. sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).

These annexes consist of a total of sheets.

3. This report contains indications relating to the following items:

- I ☒ Basis of the report
- II ☐ Priority
- III ☐ Non-establishment of opinion with regard to novelty, inventive step and industrial applicability
- IV ☐ Lack of unity of invention
- V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI ☐ Certain documents cited
- VII ☒ Certain defects in the international application
- VIII ☒ Certain observations on the international application

Date of submission of the demand 12/02/2001	Date of completion of this report 09.10.2001
Name and mailing address of the international preliminary examining authority:  European Patent Office D-80298 Munich Tel. +49 89 2399 - 0 Tx: 523656 epmu d Fax: +49 89 2399 - 4465	Authorized officer Pérez Carlón, R Telephone No. +49 89 2399 8125



INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.: PCT/EP00/06429

I. Basis of the report

1. With regard to the **elements** of the international application (*Replacement sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to this report since they do not contain amendments (Rules 70.16 and 70.17)*):

Description, pages:

1-54 as originally filed

Claims, No.:

1-10 as originally filed

2. With regard to the **language**, all the elements marked above were available or furnished to this Authority in the language in which the international application was filed, unless otherwise indicated under this item.

These elements were available or furnished to this Authority in the following language: , which is:

- ☐ the language of a translation furnished for the purposes of the international search (under Rule 23.1(b)).
- ☐ the language of publication of the international application (under Rule 48.3(b)).
- ☐ the language of a translation furnished for the purposes of international preliminary examination (under Rule 55.2 and/or 55.3).

3. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:

- ☐ contained in the international application in written form.
- ☐ filed together with the international application in computer readable form.
- ☐ furnished subsequently to this Authority in written form.
- ☐ furnished subsequently to this Authority in computer readable form.
- ☐ The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.
- ☐ The statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.

4. The amendments have resulted in the cancellation of:

- ☐ the description, pages:
- ☐ the claims, Nos.:
- ☐ the drawings, sheets:

5. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed (Rule 70.2(c)):

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/EP00/06429

(Any replacement sheet containing such amendments must be referred to under item 1 and annexed to this report.)

6. Additional observations, if necessary:

V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

1. Statement

Novelty (N)	Yes:	Claims	1-10
	No:	Claims	
Inventive step (IS)	Yes:	Claims	
	No:	Claims	1-10
Industrial applicability (IA)	Yes:	Claims	1-10
	No:	Claims	

2. Citations and explanations
see separate sheet

VII. Certain defects in the international application

The following defects in the form or contents of the international application have been noted:
s separate sheet

VIII. Certain observations on the international application

The following observations on the clarity of the claims, description, and drawings or on the question whether the claims are fully supported by the description, are made:
see separate sheet

Re Item V

Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

Reference is made to the following document:

D1: WO 98 13340

1. The application claims 3-arylsulfonyl-2-methyl propanoic acid derivatives of the formula (I) (claim 1) for use in a method of treatment(claim 7), a process for its synthesis (claim 4), pharmaceutical compositions comprising them (claim 6).
- 2 D1 (closest prior art) describes MMP-2 inhibitors that do not fall within the general formula (I) as in claim 1 of the application.

The subject-matter of claims 1-10 is therefore new, in the sense of Art. 33(2) PCT.

3. In light of the prior art, the problem to be solved can be regarded as to the provision of alternative MMP-2 inhibitors.
4. The close structural relationship between the molecules claimed and the ones described in D1 (compare for example compound 14 of D1 with compounds 8, 10 and 12 of the present application) together with its analogous activity (Ki in the order of nM and pM in both cases) cannot led to the acknowledgment of an inventive step for the subject-matter claimed. The alegued selectivity MMP-2/MMP-1 cannot be regarded as the inventiveness of the compounds of the application in the absence of evidence as to the lack of said selectivity in the compounds of the state of the art.

Moreover, only compounds 10 and 12 can be considered as having a higher activity towards MMP-2 when compared with the compounds of the art. In order to acknowledge an inventive step, the whole of the corresponding application should be inventive in the sense of Art. 33(3) PCT. To provide alternative solutions to the problem by mere (unpurposive) selection within D1 (see for example those compounds with activities in the nM range) has to be considered as obvious for a person of the art.

No inventive step in the sense of Art. 33(3) PCT is therefore acknowledged the subject-matter of claims 1-10 of the application.

5. There are no doubts about industrial applicability (Art. 33(4) PCT).

Re Item VII

Certain defects in the international application

1. The expression "and the like" at the end of lists attempts to expand the protection sought in an indeterminate way and renders the scope of the claims unclear when read in light of the description (Art. 6 PCT, the PCT Guidelines, CIII, 4.3a)

Re Item VIII

Certain observations on the international application

1. Objections as to lack of unity (Rule 13.1 PCT) could be arisen upon entry in the regional phase(s) due to the lack of inventiveness of the common structural link "X-CO-CH(CH₂-heteroatom)-CH₂-SO₂-p-C₆H₄-" between the compounds (I) of the application.
- 2.1 There is not enough support in the description (Art. 5 PCT) for claims 7-8, insofar as they refer to compounds (I) for use in a method of treatment of diseases other than those mediated by MMP-1 and MMP-2 metalloproteases.
- 2.2 There is also lack of experimental data for those inhibitors wherein R² is -S-CH₂-Ar or -O-CH₂-Ar. If a selection could in some cases be allowed within a previous application based in the differences/improvement that a structural modification might induce, generalizations from the examples to broader formulas should be regarded with special care. In this case, lack of support in the sense of Art. 5 PCT is apparent in the afore mentioned cases.

PATENT COOPERATION TREATY

PCT

INTERNATIONAL SEARCH REPORT

(PCT Article 18 and Rules 43 and 44)

Applicant's or agent's file reference FC 862/5	FOR FURTHER ACTION see Notification of Transmittal of International Search Report (Form PCT/ISA/220) as well as, where applicable, item 5 below.	
International application No. PCT/EP 00/06429	International filing date (day/month/year) 07/07/2000	(Earliest) Priority Date (day/month/year) 14/07/1999
Applicant PHARMACIA & UPJOHN SPA		

This International Search Report has been prepared by this International Searching Authority and is transmitted to the applicant according to Article 18. A copy is being transmitted to the International Bureau.

This International Search Report consists of a total of 3 sheets.



It is also accompanied by a copy of each prior art document cited in this report.

1. Basis of the report

- a. With regard to the **language**, the international search was carried out on the basis of the international application in the language in which it was filed, unless otherwise indicated under this item.



the international search was carried out on the basis of a translation of the international application furnished to this Authority (Rule 23.1(b)).

- b. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international search was carried out on the basis of the sequence listing :



contained in the international application in written form.



filed together with the international application in computer readable form.



furnished subsequently to this Authority in written form.



furnished subsequently to this Authority in computer readable form.



the statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.



the statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished

2. ☐ **Certain claims were found unsearchable** (See Box I).

3. ☐ **Unity of invention is lacking** (see Box II).

4. With regard to the **title**,



the text is approved as submitted by the applicant.



the text has been established by this Authority to read as follows:

5. With regard to the **abstract**,



the text is approved as submitted by the applicant.



the text has been established, according to Rule 38.2(b), by this Authority as it appears in Box III. The applicant may, within one month from the date of mailing of this international search report, submit comments to this Authority.

6. The figure of the **drawings** to be published with the abstract is Figure No.




as suggested by the applicant.



because the applicant failed to suggest a figure.



because this figure better characterizes the invention.



None of the figures.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 00/06429

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07C317/46 C07C323/65 C07D213/70 C07D233/74 C07D233/84
C07D235/28 C07D239/38 C07D249/12 C07D257/04 C07D277/36
C07D285/12 C07D333/34 A61K31/12 A61K31/16 A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

BEILSTEIN Data, WPI Data, EPO-Internal, PAJ, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 98 13340 A (PHARMACIA & UPJOHN) 2 April 1998 (1998-04-02) cited in the application page 2 -page 5; claims 1,11-18; examples 13-16,18-21	1,6-10
Y	WO 98 05635 A (CHIROSCIENCE) 12 February 1998 (1998-02-12) page 2 -page 4; claims 1,18-22; examples 24,56,A-D	1,6-10

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- * & * document member of the same patent family

Date of the actual completion of the international search

6 December 2000

Date of mailing of the international search report

19/12/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

English, R

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 00/06429

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61P19/02 A61P9/00 A61P25/28

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.

☐

Further documents are listed in the continuation of box C.

☒

Patent family members are listed in annex.

° Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

6 December 2000

Date of mailing of the international search report

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

English, R

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

ST/EP 00/06429

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9813340 A	02-04-1998	AU 4645997 A	17-04-1998
		BR 9712134 A	31-08-1999
		CN 1230177 A	29-09-1999
		CZ 9900923 A	14-07-1999
		EP 0929519 A	21-07-1999
		HU 0000145 A	28-07-2000
		NO 991494 A	26-05-1999
		PL 332509 A	13-09-1999
		US 5847153 A	08-12-1998
WO 9805635 A	12-02-1998	AU 3856497 A	25-02-1998
		BR 9711027 A	17-08-1999
		CN 1227540 A	01-09-1999
		CZ 9900368 A	14-07-1999
		EP 0968182 A	05-01-2000
		NO 990543 A	06-04-1999
		PL 331598 A	02-08-1999
		US 6118001 A	12-09-2000
		ZA 9707044 A	07-08-1998

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 January 2001 (25.01.2001)

PCT

(10) International Publication Number
WO 01/05756 A1

(51) International Patent Classification⁷: C07C 317/46,
323/65, C07D 213/70, 233/74, 233/84, 235/28, 239/38,
249/12, 257/04, 277/36, 285/12, 333/34, A61K 31/12,
31/16, A61P 35/00, 19/02, 9/00, 25/28

(21) International Application Number: PCT/EP00/06429

(22) International Filing Date: 7 July 2000 (07.07.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
9916562.3 14 July 1999 (14.07.1999) GB

(71) Applicant (for all designated States except US): PHAR-
MACIA & UPJOHN SpA [IT/IT]; Via Robert Koch, 1.2,
I-20152 Milan (IT).

(72) Inventors; and

(75) Inventors/Applicants (for US only): MANTEGANI,
Sergio [IT/IT]; Via C. Pisacane, 57, I-20129 Milan (IT).
ABRATE, Francesca [IT/IT]; Via Lodovico Settala,
45, I-20124 Milan (IT). BISSOLINO, Pierluigi [IT/IT];
Via Roma, 36/2, I-27020 San Giorgio Lomellina (IT).
CREMONESI, Paolo [IT/IT]; Via San Michele del Carso,

10, I-20066 Melzo (IT). PERRONE, Ettore [IT/IT]; Via
Aldo Moro, 44, I-20010 Boffalora Ticino (IT). JABES,
Daniela [IT/IT]; Via Costanza, 33, I-20146 Milan (IT).

(81) Designated States (national): AE, AL, AM, AT, AU, AZ,
BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK,
DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL,
IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU,
LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT,
RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA,
UG, US, UZ, VN, YU, ZA, ZW.


(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- With international search report.
- Before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments.

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: 3-ARYLSULFONYL-2-(SUBSTITUTED METHYL)PROPANOIC ACID DERIVATES AS MATRIX METALLOPRO-
TEINASE INHIBITORS



(I)

(57) Abstract: Compounds which
are 3-arylsulfonyl-2-methyl propanoic
acid derivatives of formula (I):
wherein X is HO-NH- or HO-, R₁ is
selected from phenyl, 4-chlorophenyl,
4-fluorophenyl, 4-cyanophenyl,
benzamido (i.e., -NH-CO-Ph) and
benzamido substituted on the terminal
phenyl ring by C₁-C₄ alkyl, fluoro,
chloro, cyano or C₁₋₄ alkoxy; R₂ is
selected from (a) -S-Ar or -S-CH₂-Ar
wherein Ar is an aromatic moiety;
(b) -O-Ar wherein Ar is as defined
above; (c) -S-Het or -S-CH₂-Het
wherein Het is a heterocyclic ring;

and (d) 2,5-dioxo-1-imidazolidinyl or 2,4-dioxo-1-imidazolyl; and the pharmaceutically acceptable salts thereof; have potent
and selective inhibitory activity against matrix metalloproteinases (MMPs) and can thus be used in the treatment and prevention
of diseases mediated by MMPs.

3-ARYLSULFONYL-2-(SUBSTITUTED METHYL)PROPANOIC ACID DERIVATES AS MATRIX METALLOPROTEINASE INHIBITORS

FIELD OF THE INVENTION

5 The present invention relates to some β -sulfonyl hydroxamic and carboxylic acids displaying potent and selective inhibitory activity against some Matrix MetalloProteinases (MMPs), to their use for the treatment of diseases in which the proteolytic activity of these MMPs is involved, to methods for their preparation, and to pharmaceutical compositions containing them.

BACKGROUND OF THE INVENTION


10 Under physiological conditions MMPs play an important role in normal tissue turnover and development. However, increased MMP activity and/or impairment of the physiological MMP inhibitors has been associated with a variety of pathological conditions such as cancer, rheumatoid arthritis, osteoarthritis, osteoporosis, multiple
15 sclerosis, stroke, arteriosclerosis, restenosis and age-related macular degeneration. In cancer, the contribution of MMPs is manifold and not limited to destruction of the extracellular matrix required for local and distal invasion. The proteolytic action of MMPs generates active matrix protein fragments, and influence the release, activation and bioavailability of growth factors. In addition, MMPs are involved in cell migration and in
20 the processing or shedding of cell surface proteins. It is now believed that the proteolytic action of MMPs plays a very important role in tumor growth, apoptosis and angiogenesis (see Noel, A. et al., 'Emerging Roles for Proteinases in Cancer', Invasion Metastasis 17:221-239, 1997). Among the several MMPs known to date, gelatinase-A (MMP-2) is currently reputed as the main target in cancer.

25 Potent MMP inhibitors (MMPIs) were disclosed three decades ago (e.g., see Dickens, J. P. et al., U.S. Patent 4,599,361), but only recently some of them entered clinical trials, and none has arrived at the market yet. Questions remain concerning their specificity, bioavailability, and potential long-term toxicity (see Hodgson, J., 'Remodeling MMPIs', Biotechnology 13:554-557, 1995), especially because they are primarily intended
30 (in arthritis and cancer, at least) for prolonged therapy. Marimastat, the most advanced

MMPI in the clinic, and other compounds currently investigated in patients such as Novartis CGS27023A and Agouron AG3340, have been reported to cause a musculoskeletal syndrome characterized by joint pain and stiffness, which may be severe and dose-limiting. Although the cause of this toxicity is not well understood at this moment, it is thought to arise from an impairment in the normal tissue remodeling that is governed by one or more of the MMPs, primarily fibroblast collagenase (MMP-1). We have extended these observations by developing an animal model of joint toxicity, involving treatment of rats for 10 days with MMPis at exposures close to that required for antitumor efficacy in mice (human prostate DU-145 model). In this rat model, MMPis characterized by poor selectivity (i.e., comparable potency against MMP-2 and MMP-1), such as marimastat and Roche Ro31-9790, elicited histological alteration of stifle joints (hypertrophic fibrosis of ligaments, interstitial hypertrophic fibrosis of skeletal muscles, hypertrophic fibroplasia of the periostium and synovium, chondrodysplasia, and decreased endochondrial ossification of the ephyseal plate); in some cases, clinical symptoms (abnormal posture, reluctance to move, abnormal gait) were also observed. Thus, there is a strong need for better MMPis, especially as far as the properties referred to above (specificity, bioavailability, long-term toxicity) are concerned. The compounds of the present invention are intended to provide these advantages.

INFORMATION DISCLOSURE

A previous disclosure by the Pharmacia & Upjohn Company, PCT Patent Application WO 98/13340 (US Patent Number 5,847,153), provides a broad range of β -sulfonyl hydroxamic acid derivatives, their preparation, and their activity against various enzymes of the MMP family, predominantly stromelysin and gelatinase. These compounds are generically described by the formula (Ia):




wherein R_1 is C_{4-12} alkyl, C_{4-12} alkenyl, C_{4-12} alkynyl, $-(CH_2)_h-C_{3-8}$ cycloalkyl, $-(CH_2)_h$ -aryl, or $-(CH_2)_h$ -het; R_2 is C_{1-12} alkyl, C_{2-12} alkenyl, C_{2-12} alkynyl, $-(CH_2)_h-C_{3-8}$ cycloalkyl, $-(CH_2)_h-C_{3-8}$ cycloalkenyl, $-(CH_2)_h$ -aryl, $-(CH_2)_h$ -het, $-(CH_2)_h$ -Q, $-(CH_2)_l$ -X- R_4 , or $-(CH_2)_l$ -CHR₅R₆.

5 We have now found that the combination of few particular groups R_1 and R_2 in a molecule of formula (Ia) provides distinct advantages over the other compounds of this class of MMPis. Said combination, which is an object of the present invention, is associated with outstanding potency (K_i in the picomolar range) against the gelatinases, in particular MMP-2, and relatively weak activity against fibroblast collagenase (MMP-1).
 10 Further, we have found that the same combination confers good potency (low nanoM range) and even higher selectivity for MMP-2 to the corresponding carboxylates, which were previously described as intermediates but not claimed as such. Finally, we have found that the same combination provides advantages in the pharmacokinetic parameters, in particular lower clearance and higher oral bioavailability, and that antitumor efficacy in
 15 rodents can be achieved in the absence of joint toxicity.

DESCRIPTION OF THE INVENTION

The present invention provides 3-arylsulfonyl-2-methyl propanoic acid derivatives of formula (I):



20

or pharmaceutically acceptable salts thereof wherein:

X is either HO-NH- (hydroxamic acids) or HO- (carboxylic acids);

R_1 is selected from phenyl, 4-chlorophenyl, 4-fluorophenyl, 4-cyanophenyl, benzamido (i.e., -NH-CO-Ph) or benzamido substituted on the terminal Ph ring by C_{1-4} alkyl, fluoro, chloro, cyano, C_{1-4} alkoxy;

25

R₂ is selected from:

(a) -S-Ar or -S-CH₂-Ar, wherein Ar is a monocarbocyclic or bicarbocyclic aromatic moiety, preferably phenyl, naphthyl and biphenyl, which is either unsubstituted or substituted with one or two substituents selected from the group consisting of C₁₋₄ alkyl, phenyl, benzyl, C₁₋₄ alkoxy, fluoro, chloro, bromo, nitro, cyano, hydroxy, amino, dimethylamino, acetamido, methylthio and acetyl;

(b) -O-Ar, wherein Ar is as defined above;


(c) -S-Het or -S-CH₂-Het, wherein Het is a heterocyclic ring selected from the group consisting of pyridine, pyrimidine, pyridazine, pyrazine, 1,2,5-triazine, imidazole, thiophene, furan, pyrrole, pyrazole, 1,3-thiazole, 1,3-oxazole, 1,2,3-triazole, 1,2,4-triazole, 1,3,4-thiadiazole, 1,3,4-oxadiazole, 1,2,3,4-tetrazole, quinoline, isoquinoline, indole, 1,3-benzoxazole, 1,3-benzothiazole, benzimidazole, [1,3]oxazolo[4,5-b]pyridine, [1,3]thiazolo[4,5-b]pyridine, [1,2,3,4]tetrazolo[1,5-b]pyridazine and purine, and wherein said Het group can be substituted with one to three substituents selected from the group consisting of C₁₋₄ alkyl, phenyl, pyridyl, benzyl, C₁₋₄ alkoxy, methylthio, fluoro, chloro, nitro, cyano, hydroxy, oxo, amino, methylamino, dimethylamino, 2-dimethylaminoethyl, acetamido and acetyl;

(d) 2,5-dioxo-1-imidazolidinyl or 2,4-dioxo-1-imidazolidinyl, which can be substituted at the carbon atom with one or two methyl groups, or with C₂₋₄ linear or branched alkyl, phenyl, benzyl or hydroxymethyl, and at the nitrogen atom with C₁₋₄ linear or branched alkyl.

The compounds of the present invention can be converted to pharmaceutically acceptable salts, where appropriate, according to conventional methods. The term 'pharmaceutically acceptable salts' refers to acid addition salts useful for administering the compounds of this invention and includes hydrochloride, sulfate, phosphate, acetate, propionate, lactate, mesylate, maleate, malate, succinate, tartrate, citrate, 2-hydroxyethyl sulfonate, fumarate and the like. Alternatively, the compounds of this invention, especially when X in formula (I) is hydroxy, may form metal salts, such as sodium, potassium, calcium or magnesium, or ammonium salts, or salts with an appropriate organic amine or amino acid such as arginine, procaine and the like. Any salt may be either in the anhydrous or hydrated form, or may be solvated with acceptable solvents, such as ethanol.

The present invention also provides pharmaceutical compositions which comprise, as active ingredient, a compound of formula (I) or a solvate, hydrate, or pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier, diluent or excipient.

The compounds of this invention contain a chiral center at the α -position of the hydroxamic or carboxylic acid, and as such they may be obtained as any of the two separate enantiomers or as a racemic mixture of both. Preferred compounds of this invention are either the racemates or individual isomers having the configuration depicted in the formula (I') herebelow:



10

wherein X, R_1 and R_2 are as defined above. According to the Cahn-Ingold-Prelog rule, these preferred isomer are denoted as (*R*) when R_2 is $-\text{O-Ar}$, 2,5-dioxoimidazolidin-1-yl or 2,4-dioxoimidazolidin-1-yl, and as (*S*) when R_2 is $-\text{S-Ar}$, S-Het, $-\text{S-CH}_2\text{-Ar}$ or $-\text{S-CH}_2\text{-Het}$. In addition, depending on the substituents, additional chiral centers or other isomeric forms may be present, and this invention embraces all possible stereoisomers and geometrical forms.

Particularly preferred compounds of this invention are as follows:

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfonyl)methyl]propanoic acid;

(2*S*)-3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfonyl)methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(phenylsulfonyl)methyl]-propanamide;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[4-hydroxyphenyl)sulfonyl]methyl]propanoic acid;

25

- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(4-hydroxyphenyl)sulfanyl]-methyl]propanamide;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-(phenoxymethyl)propanoic acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-(phenoxymethyl)propanamide;
- 5 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2-pyridinylsulfanyl)methyl]propanoic acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(2-pyridinylsulfanyl)methyl]-propanamide;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2-pyridinylmethyl)sulfanyl]methyl]-10 propanoic acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(2-pyridinylmethyl)sulfanyl]-methyl]propanamide;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(5-methyl-1,3,4-thiadiazol-2-yl)sulfanyl]-methyl]propanoic acid;
- 15 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(5-methyl-1,3,4-thiadiazol-2-yl)sulfanyl]methyl]propanamide;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(5-amino-1,3,4-thiadiazol-2-yl)-sulfanyl]-methyl]propanoic acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(1-methyl-1H-imidazol-2-yl)sulfanyl]-20 methyl]propanoic acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2-pyrimidinylsulfanyl)methyl]propanoic acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(1-methyl-1H-tetrazol-5-yl)sulfanyl]methyl]-25 propanoic acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(1-benzyl-1H-tetrazol-5-yl)sulfanyl]methyl]-propanoic acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(1H-benzimidazol-2-yl)sulfanyl]methyl]-30 propanoic acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(1,3-thiazol-2-yl)sulfanyl]methyl]propanoic

- acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[(2-methyl-2H-1,2,3-triazol-4-yl)sulfanyl]-methyl]propanoic acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[(5-phenyl-1,3,4-oxadiazol-2-yl)sulfanyl]-methyl]propanoic acid;
- 5 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[1-(3-pyridinyl)-1H-imidazol-2-yl)sulfanyl]-methyl]propanoic acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[(4-methyl-1,3-oxazol-2-yl)sulfanyl]methyl]-propanoic acid;
- 10 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[1,3]oxazolo[4,5-b]pyrimidin-2-yl-sulfanyl)-methyl]propanoic acid;
- 15 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[1,2,3,4]tetrazolo[1,5-b]pyridazin-6-yl-sulfanyl)methyl]propanoic acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[(2,6-dimethyl-4-pyrimidinyl)sulfanyl]-methyl]propanoic acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[(4,6-dimethyl-2-pyrimidinyl)sulfanyl]-methyl]propanoic acid;
- 20 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[(4-hydroxy-6-methyl-2-pyrimidinyl)-sulfanyl]methyl]propanoic acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[(4,6-dihydroxy-2-pyrimidinyl)sulfanyl]-methyl]propanoic acid;
- 25 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(9H-purin-6-ylsulfanyl)methyl]propanoic acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-(3-fluorophenoxymethyl)propanoic acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2-acetylaminophenoxy)methyl]propanoic acid;
- 30 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(4-acetylaminophenoxy)methyl]propanoic acid;


- 3-([1,1'-Biphenyl]-4-ylsulfonyl)-2-[(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)methyl]-
propanoic acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3,4,4-trimethyl-2,5-dioxo-1-
imidazolidinyl)-
5 methyl]propanoic acid;
- (2R)-3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3,4,4-trimethyl-2,5-dioxo-1-
imidazolidinyl)methyl]propanoic acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(3,4,4-trimethyl-2,5-dioxo-1-
imidazolidinyl)methyl]propanamide;
- 10 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(4,4-dimethyl-2,5-dioxo-1-imidazolidinyl)-
methyl]propanoic acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(4,4-dimethyl-2,5-dioxo-1-
imidazolidinyl)methyl]propanamide;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2,5-dioxo-1-
15 imidazolidinyl)methyl]propanoic
acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(2,5-dioxo-1-imidazolidinyl)-
methyl]propanamide;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3-methyl-2,5-dioxo-1-imidazolidinyl)-
20 methyl]propanoic acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3-ethyl-2,5-dioxo-1-
imidazolidinyl)methyl]-
propanoic acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3-butyl-2,5-dioxo-1-
25 imidazolidinyl)methyl]-
propanoic acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(3-butyl-2,5-dioxo-1-
imidazolidinyl)methyl]propanamide;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3-butyl-2,4-dioxo-1-
30 imidazolidinyl)methyl]-
propanoic acid;

- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(3-butyl-2,4-dioxo-1-imidazolidinyl)methyl]propanamide;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(5-isopropyl-2,4-dioxo-1-imidazolidinyl)methyl]propanoic acid;
- 5 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(5-hydroxymethyl-2,4-dioxo-1-imidazolidinyl)methyl]propanoic acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(5-hydroxymethyl-3-methyl-2,4-dioxo-1-imidazolidinyl)methyl]propanoic acid;
- 3-[(4'-Fluoro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfanyl)methyl]propanoic acid;
- 10 3-[(4'-Fluoro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(phenylsulfanyl)methyl]-propanamide;
- 3-[(4'-Fluoro[1,1'-biphenyl]-4-yl)sulfonyl]-2-(phenoxymethyl)propanoic acid;
- 3-[(4'-Fluoro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-(phenoxymethyl)propanamide;
- 3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfanyl)methyl]propanoic acid;
- 15 3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(phenylsulfanyl)methyl]-propanamide;
- 3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-(phenoxymethyl)propanoic acid;
- 3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-(phenoxymethyl)propanamide;
- 3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-(3-fluorophenoxymethyl)propanoic acid;
- 20 3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2-acetylamino)phenoxymethyl]propanoic acid;
- 3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2-pyridinylsulfanyl)methyl]propanoic acid;
- 3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2-pyridinylmethyl)sulfanyl)methyl]propanoic acid;
- 25 3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3-butyl-2,5-dioxo-1-imidazolidinyl)methyl]-propanoic acid;
- 3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(3-butyl-2,5-dioxo-1-imidazolidinyl)methyl]propanamide;
- 30 3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(5-hydroxymethyl-2,4-dioxo-1-imidazolidinyl)methyl]propanoic acid;

- 3-[[4-(Benzoylamino)phenyl]sulfonyl]-2-[(phenylsulfanyl)methyl]propanoic acid;
3-[[4-(Benzoylamino)phenyl]sulfonyl]-N-hydroxy-2-[(phenylsulfanyl)methyl]-
propanamide;
3-[[4-(Benzoylamino)phenyl]sulfonyl]-2-[[2-(pyridinylmethyl)sulfanyl]methyl]propanoic
5 acid;
3-[[4-(Benzoylamino)phenyl]sulfonyl]-2-[(2-thienylsulfanyl)methyl]propanoic acid;
3-[[4-(Benzoylamino)phenyl]sulfonyl]-2-(phenoxymethyl)propanoic acid;
3-[[4-[(4-Methylbenzoyl)amino]phenyl]sulfonyl]-2-[(phenylsulfanyl)methyl]propanoic
acid;
10 3-[[4-[(4-Methylbenzoyl)amino]phenyl]sulfonyl]-N-hydroxy-2-[(phenylsulfanyl)methyl]-
propanamide;
3-[[4-[(4-Chlorobenzoyl)amino]phenyl]sulfonyl]-2-[(phenylsulfanyl)methyl]propanoic
acid;
3-[[4-[(4-Chlorobenzoyl)amino]phenyl]sulfonyl]-N-hydroxy-2-[(phenylsulfanyl)methyl]-
15 propanamide;
3-[[4-[(4-Chlorobenzoyl)amino]phenyl]sulfonyl]-2-(phenoxymethyl)propanoic acid;
3-[[4-[(4-Chlorobenzoyl)amino]phenyl]sulfonyl]-N-hydroxy-2-(phenoxymethyl)-
propanamide;
3-[[4-[(4-Chlorobenzoyl)amino]phenyl]sulfonyl]-2-(3-fluorophenoxymethyl)propanoic
20 acid;
3-[[4-[(4-Chlorobenzoyl)amino]phenyl]sulfonyl]-2-[[2-(pyridinylmethyl)sulfanyl]methyl]-
propanoic acid;
3-[[4-[(4-Chlorobenzoyl)amino]phenyl]sulfonyl]-2-[[1-methyl-1H-tetrazol-5-yl)sulfanyl]-
methyl]propanoic acid;
25 3-[[4-[(4-Chlorobenzoyl)amino]phenyl]sulfonyl]-2-[(5-hydroxymethyl-2,4-dioxo-1-
imidazolidinyl)methyl]propanoic acid;
3-[[4-[(4-Cyanobenzoyl)amino]phenyl]sulfonyl]-2-[(phenylsulfanyl)methyl]propanoic
acid;
3-[[4-[(4-Cyanobenzoyl)amino]phenyl]sulfonyl]-N-hydroxy-2-[(phenylsulfanyl)methyl]-
30 propanamide;
3-[[4-[(4-Cyanobenzoyl)amino]phenyl]sulfonyl]-2-[[1-methyl-1H-tetrazol-5-yl)sulfanyl]-

- methyl]propanoic acid;
 3-[[4-[(4-Cyanobenzoyl)amino]phenyl]sulfonyl]-2-(phenoxymethyl)propanoic acid;
 3-[[4-[(4-Cyanobenzoyl)amino]phenyl]sulfonyl]-N-hydroxy-2-(phenoxymethyl)-
 propanamide;
 5 3-[[4-[(4-Cyanobenzoyl)amino]phenyl]sulfonyl]-2-(3-fluorophenoxymethyl)propanoic
 acid;
 3-[[4-[(4-Cyanobenzoyl)amino]phenyl]sulfonyl]-N-hydroxy-2-(3-fluorophenoxymethyl)-
 propanamide;
 3-[[4-[(4-Cyanobenzoyl)amino]phenyl]sulfonyl]-2-[(2-acetylamino)phenoxymethyl]-
 10 propanoic acid;
 3-[[4-(4-Propylbenzoylamino)phenyl]sulfonyl]-2-[[1-methyl-1H-tetrazol-5-yl)sulfanyl]-
 methyl]propanoic acid;
 3-[[4-[(4-Methoxybenzoyl)amino]phenyl]sulfonyl]-2-(phenoxymethyl)propanoic acid;
 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[2-(hydroxymethyl)phenoxy]methyl]-
 15 propanoic acid.

The compounds of this invention can be prepared by a process starting from a compound of formula 4:



- 20 wherein R is H or the residue of a carboxylic acid ester, R_1 and R_2 are as defined
 above and n is 0 or 2, which process comprises:

- (a) hydrolysing a said compound of formula 4 in which R is the residue
 of a carboxylic and ester to give a compound of formula (I) in which X is HO-; or
 (b) hydrolysing and oxidising, in either order, a said compound of
 25 formula 4 in which n is 0 and R is the residue of a carboxylic acid ester, to give a
 compound of formula (I) in which X is HO-; or


(c) activating a said compound of formula 4 wherein R is H and n is 2 to form an activated carboxy group, coupling the activated carboxy group with hydroxylamine or an O-protected derivative thereof and, if necessary, deprotecting the hydroxamic group to give a compound of formula (I) wherein X is -NHOH; or

5 (d) submitting a said compound of formula 4 wherein R is H and n is zero to a sequence of reactions comprising oxidation at the sulphur atom, activation of the carboxy group, condensation of the activated carboxy group with hydroxylamine or an O-protected derivative thereof and, if necessary, deprotection of the hydroxamic group to form a compound of formula (I) wherein X is -NHOH, the oxidation step being conducted
10 either before the activation step or after the condensation step; and/or

(e) if desired, converting a resulting compound of formula (I) into another compound of formula (I); and/or converting a free compound into a pharmaceutically acceptable salt thereof; and/or converting a salt into a free compound.


The compound of formula 4 may be obtained by

15 (a) subjecting a compound of formula 2:




wherein R, R₁ and n are as defined above, to conjugate addition by treatment with a compound of formula R₂H wherein R₂ is as defined above; or

20 (b) treating a compound of formula 3:



wherein R and R₂ are as defined above, with a thiol of formula:



to obtain a compound of formula 4 in which n is zero.

These processes are depicted in the following Scheme I, where R₁ and R₂ are the groups as defined previously, R is either a hydrogen atom or the residue of a carboxylic ester such as methyl, ethyl, n-butyl, tert-butyl, benzyl or benzhydryl, and n is either zero or 2. Compound 1, i.e. 2-(bromomethyl)acrylic acid or an ester thereof, is reacted with the appropriate aryl-thiol in a suitable solvent such as toluene or dimethylformamide (DMF) in the absence or presence of a suitable base such as sodium bicarbonate or triethylamine (TEA), at ambient temperature or under heating up to reflux temperature, to afford a compound 2 wherein n is zero. The corresponding sulfone (i.e., a compound 2 wherein n is 2) is obtained, if desired, by subsequent reaction with an oxidant such as metachloroperbenzoic acid (MCPBA) in a suitable solvent such as dichloromethane (DCM) at 0 °C or ambient temperature, or using hydrogen peroxide in acidic acid; preferably, this reaction is performed with potassium peroxymonosulfate (Oxone[®]) in an organic solvent, preferably DMF, or in a mixture of water and water-miscible organic solvents such as methanol, DMF or the combination of two, at ambient temperature or under moderate heating. Alternatively, a compound 2 wherein n is 2 can be obtained from compound 1 directly, by reaction with the appropriate arylsulfinate. Preferably, the sulfinate is obtained and used as the sodium salt, or as the salt with a tertiary organic amine such as N-methylmorpholine (NMM). The sodium arylsulfinate can be conveniently obtained by reaction of the appropriate aryl-sulfonyl chloride with sodium iodide in acetone, preferably at ambient temperature, and it is isolated before reaction with compound 1, which takes place in a suitable solvent such as toluene in the absence or presence of a suitable base such as sodium bicarbonate, at a temperature ranging from 25 °C to reflux. The salt with NMM is obtained by reaction of the arylsulfonyl chloride with 4-thiocresol and NMM in DCM, preferably at a temperature ranging from -50 to -78 °C, and it is reacted in situ with compound 1, then raising the temperature up to 25 °C or reflux.

A compound 4 can be obtained from the corresponding compound 2 by conjugate addition of a reagent of formula R_2H , or a salt thereof, where R_2 is as defined previously. Thus, when R_2 is a group $-S-Ar$, $-S-Het$, $-S-CH_2-Ar$ or $-S-CH_2-Het$, the reagent is the corresponding thiol, or a salt thereof; when R_2 is $-O-Ar$, the reagent is the corresponding aromatic carbinol; when R_2 is 2,5-dioxo-1-imidazolidinyl or a substituted derivative thereof, the reagent is the corresponding 2,4-imidazolidinedione. Said conversion from a compound 2 into a compound 4 can be accomplished, under identical or similar conditions, either when n in 2 is zero or it is 2; however, the preparation of compounds of formula (I) wherein R_2 is sulfanyl is better performed from compounds 2 where n is 2, because otherwise it is difficult to discriminate between the two sulfur atoms in the following oxidative step. Typically, the conjugate addition of said mercaptans and aromatic carbinols to a compound of formula 2 is accomplished using the reagents neat, or in a suitable organic solvent such as DCM, DMF, toluene, tetrahydrofuran (THF), acetonitrile (MeCN), methanol or ethanol, at temperatures ranging from ambient to reflux, in the absence or presence of a suitable base such as sodium bicarbonate, sodium carbonate, TEA, sodium methoxide or sodium ethoxide, preferably under a nitrogen blanket. When the reagent is 2,4-imidazolidinedione or a substituted derivative thereof, the conjugate addition to a compound 2 is preferably performed in toluene or DMF in the presence of a suitable base such as sodium bicarbonate, TEA, NMM and the like, at a temperature ranging from ambient to reflux, preferably between 50 and 110 °C; alternatively, a preformed salt of said reagent can be used, especially sodium, potassium and lithium salts, which can be obtained by treatment with suitable strong bases, especially sodium hydride, potassium hexamethyldisilazide, potassium tert-butoxide and butyllithium, in a suitable organic solvent.

When a compound 4 where n is zero is obtained by the methods above, it may be converted, if desired, into the corresponding compound where n is 2 by oxidation, under the same conditions described previously for the oxidation of a compound 2 where n is zero to a compound where n is 2.

Alternatively, a compound 4 can be obtained from a compound 1 through the intermediacy of a compound 3. This alternative route provides advantages when the reagent R_2H is reluctant to the conjugate addition to a compound 2, e.g. when R_2H is 2,5-

dioxo-1-imidazolidine, and particularly when it is 4,4-dimethyl-2,5-dioxo-1-imidazolidine and the N-alkyl derivatives thereof. In the first step, said reagent R_2H is condensed with the bromomethylacrylate 1 to provide 3. Typically, the reaction is performed in a solvent such as DMF or toluene, at temperatures ranging from 50 °C to reflux, and preferably in the presence of sodium bicarbonate. In the following step, conjugate addition of the suitable aryl-thiol to the intermediate 3 provides a compound 4 where n is zero. This reaction can be performed under the same conditions described above for reaction of 2 with a mercaptane R_2H to provide 4.

Compounds 4 where R is a hydrogen atom and n is 2 are specifically represented by structure 5 and as such are a first object of this invention, i.e. the compounds of formula (I) wherein X is hydroxy (the 'carboxylates'). Compounds 4 where R is the residue of a carboxylic ester, as defined previously, and n is 2, are converted to said final compounds 5 by hydrolysis, according to the general methods known in the art such as using 6N HCl in acetic acid and refluxing for 10 to 20 hours or using iodotrimethylsilane in chloroform. When the R group is benzyl and R_2 is not a sulfanyl residue, a preferred method to obtain the free acid is hydrogenolysis, especially in the presence of a palladium catalyst such as Pd on charcoal, in an inert organic solvent such as ethanol or DMF or the like, typically at room temperature and under atmospheric pressure or moderate pressure. When the R group is tert-butyl, a preferred method for obtaining the free acid is reaction with HCl or trifluoroacetic acid, in inert organic solvents such as DCM, THF, MeCN and the like, optionally in admixture with water, methanol or ethanol, at temperatures ranging from -20 to +40 °C.


Compounds 4 where R is the residue of a carboxylic ester and n is zero are converted to compounds 5 by a two-step procedure involving oxidation and hydrolysis, or hydrolysis and oxidation, as preferred. Conditions for oxidation and hydrolysis are the same as described above.

Compounds 4 where R is a hydrogen atom and n is 2, which have structure 5, are converted to compounds 6, which specifically represent the second object of this invention, i.e. structure (I) wherein X is -NHOH (the 'hydroxamates'), by activation of the acid and coupling with hydroxylamine or an O-protected derivative thereof. This reaction can be performed by several routes well known to those skilled in the art. For example, the

acid can be activated as its chloride or a mixed anhydride. A preferred method for activation as the acid chloride is reaction with oxalyl chloride, typically in DCM containing DMF as a catalyst, at 0 °C to ambient temperature. A preferred method for activation as a mixed anhydride is reaction with a chlorocarbonate such as chloroethylformate in anhydrous THF or a similar compatible solvent, in the presence of a tertiary amine such as TEA, NMM and the like. Alternatively, the acid can be activated by a carbodiimide condensing agent such as dicyclohexylcarbodiimide (DCC) or 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (WSCDI, water soluble carbodiimide), preferably in the presence of 1-hydroxybenzotriazole (HOBt) in an inert organic solvent such as DMF at 0 °C to room temperature. Another method for the activation of the acids of formula 4 where R is hydrogen is reaction with benzotriazolyl-1-yloxy-tris(dimethylamino)phosphonium hexafluorophosphate (BOP reagent) in a suitable organic solvent such as MeCN, THF or DCM in the presence of a tertiary amine such as TEA or NMM. A further method is reaction with O-benzotriazolyl-N,N,N',N'-tetramethyluronium hexafluorophosphate (HBTU) or O-benzotriazolyl-N,N,N',N'-tetramethyluronium tetrafluoroborate (TBTU), under conditions similar to those described for the BOP reagent. When the activation is performed with oxalyl chloride, the acid chloride is usually isolated crude before reaction with the hydroxylamine reagent; when with chloroethylformate, EDC or WSCDI, the hydroxylamine reagent is preferably added after about 1 hour at room temperature; when with the BOP reagent, HBTU or TBTU, the hydroxylamine reagent is O-protected and it is preferably added from the beginning. Suitable O-protected hydroxylamines are O-(benxyl)benzylhydroxylamine, O-(tert-butyl)hydroxylamine, O-(tetrahydropyranyl)-hydroxylamine, O-(trimethylsilyl)hydroxylamine and O-(tert-butyldimethylsilyl)-hydroxylamine, the first two being preferably used as their hydrochlorides and freed in situ by a tertiary amine such as TEA or NMM. Alternatively, when activation of the acid is performed with oxalyl chloride or WSCDI, the unprotected hydroxylamine (e.g., as the commercial 50% solution in water) or the hydrochloride thereof can be used, either in the organic solvents specified above or after dissolution in water or in a mixture of water and THF. Suitable conditions for the condensation step vary according to the activation method. Typically, when activation is by the acid chloride method, the reaction is fast and is complete within 1-5

hours at temperatures ranging from 0 °C to 25 °C; otherwise it can be slower, usually requiring stirring overnight at ambient temperature or even 1-5 days. When an O-protected hydroxylamine is used in the condensation, the reaction affords O-protected hydroxamate derivatives, and the desired free-hydroxamates 6 are obtained after removal of the protecting group. In particular, when O-(benzyl)hydroxylamine is used, deprotection is performed by hydrogenolysis, preferably with a palladium catalyst such as 5% Pd on charcoal, 5% Pd on BaSO₄, or palladium hydroxide on carbon (Pearlman's catalyst), typically in methanol or ethanol. When O-(tert-butyl)hydroxylamine is used, the free hydroxamate can be obtained after treatment with neat or aqueous trifluoroacetic acid, or with HCl in DCM. When O-(tetrahydropyranyl)hydroxylamine is used, the free hydroxamate is rapidly obtained by reaction with 4N aqueous HCl in suitable organic solvents or mixtures thereof, e.g. dioxane and methanol. When O-(trimethylsilyl)hydroxylamine is used, deprotection is usually achieved during work up, e.g. by partitioning of the condensation mixture between ethyl acetate and diluted hydrochloric acid. When O-(tert-butyltrimethylsilyl)hydroxylamine is used, deprotection can be achieved by HF or fluoride salts, preferably with tetrabutylammonium fluoride, in a suitable organic solvent such as THF, at temperatures ranging from -10 to +25 °C.

Compounds 4 where R is a hydrogen atom and n is zero are converted to compounds 6 by a sequence involving oxidation at the sulfur atom, activation of the carboxy group, condensation of the activated carboxy group with hydroxylamine or an O-protected derivative thereof and, when required, unmasking of the free hydroxamic group. The oxidation can be performed first, i.e. on said compounds of formula 4, as described previously for the oxidation of a compound of formula 2 where n is zero to the corresponding compound where n is 2. Alternatively, the oxidation can be performed at a later stage, i.e. on the O-protected or free hydroxamic acid, in which case a preferred oxidant is oxone, in a suitable organic solvent such as DMF, mixture of water and one or two water miscible organic solvents, such as methanol, ethanol, DMF and THF, at a temperature ranging from 5 to 25 °C. Conditions for the other steps of the sequence, i.e. activation of the carboxy group, coupling with hydroxylamine or a protected derivative thereof and, if required, deprotection, are the same as described previously.

SCHEME I


5 In some cases, minor modifications of the general procedure indicated in Scheme 1 can be used to obtain compounds 4. For example, conversion of a compound 2 into a compound 4 wherein R₂ is a sulfanyl residue can be achieved in two steps, the first involving addition of HBr to a compound 2 to obtain the bromomethyl derivative, which is then reacted with the thiol which R₂H can represent. This two-step procedure can be

10 convenient for less reactive thiols, in particular when R₂ is -S-CH₂-Ar or -S-CH₂-Het. A preferred solvent for the addition of HBr to a compound 2 is acetic acid, usually at ambient temperature. The thiol is preferably reacted with said bromomethyl intermediate in the presence of an organic or inorganic base such as TEA, NMM, sodium bicarbonate,

potassium carbonate and the like. Less reactive thiols can be advantageously reacted after conversion to salts thereof, e.g. by treatment with potassium hexamethyldisilazide at low temperature in a suitable organic solvent such as THF, and reacted in situ with the compound 2. The bromomethyl intermediate, or the corresponding carbinol, can also be
5 used to obtain compounds 4 wherein R_2 is 2,5-dioxo-1-imidazolidinyl. In particular, the bromomethyl derivative can be reacted with 2,4-imidazolidinedione, or a derivative thereof, under conditions similar to that described above for reaction with a thiol, and the carbinol can be condensed with the same reagent under Mitsunobu conditions, i.e. in the presence of triphenylphosphine and diethyl azodicarboxylate in DMF at temperatures
10 ranging from 0 to 25 °C.

The compounds of formula (I) wherein R_2 is 2,4-dioxo-1-imidazolidinyl can be obtained by the variant described in Scheme II below. As therein outlined, an aminoacid H_2N-CHR_3-COOH where R_3 is a hydrogen atom, C_{1-4} straight or branched alkyl or benzyl, or a salt or an ester thereof, especially the methyl or ethyl esters, is condensed with
15 compound 1, as defined in Scheme I, to obtain a compound 7 where R , R_3 are as above defined, and R' is either a hydrogen atom or methyl or ethyl. This reaction is carried out in a suitable organic solvent, with or without a base, at temperatures ranging from 0 to 50 °C, preferably around ambient temperature. A preferred solvent is DMF; preferred bases, which can be present in amounts variable to a catalytic one to 3 molar equivalents, are
20 tertiary amines, such as TEA, NMM, Hunig's base and the like. The obtained compound 7 is then reacted with sodium or potassium cyanate, or with an isocyanate of formula R_4-NCO where R_4 is C_{1-4} alkyl, phenyl or benzyl, to obtain an intermediate urea, which is cyclized to a compound 8 where R , R_3 and R_4 are as defined above, in a suitable organic solvent such as THF, dioxane, MeCN, DMF and the like, at temperatures ranging from
25 about 20 °C to reflux. Said compound 8 is then reacted with an aryl thiol of formula $R_1-C_6H_4-SH$ to obtain a compound 9 where R_1 is as above defined and n is zero. Typically, the conjugate addition of said aryl thiol to a compound 8 is accomplished in a suitable organic solvent such as DCM, DMF, toluene, tetrahydrofuran (THF), acetonitrile (MeCN), methanol or ethanol, at temperatures ranging from ambient to reflux, in the absence or
30 presence of a suitable base such as sodium bicarbonate, sodium carbonate, TEA, sodium methoxide or sodium ethoxide, preferably under a nitrogen blanket. Conversion of a

compound 9 where n is zero into a compound 9 where n is 2 is accomplished under the conditions previously described in Scheme I, preferably with Oxone[®] in an organic solvent such as DMF or in a mixture of water and water-miscible organic solvent such as methanol, DMF or the combination of two, at ambient temperature or under moderate heating. Alternatively, the same synthetic sequence is carried out on a compound 2, as defined in Scheme I, to obtain the same compound 9 *via* 10. Compounds 9 where n is two and R is a hydrogen atom are the desired compounds of formula (I) where X is HO- and R₂ is 2,4-dioxoimidazolidin-1-yl. The corresponding compounds of formula (I) where X is HONH- are obtained, if desired, as described previously (Scheme I) for the conversion of a compound 5 into a compound 6.

SCHEME II

When required or convenient, chemical modifications of the group R_1 or R_2 in intermediates 2, 3, 4 and 9 can be performed in the above processes to obtain the final compounds of formula (I). For example, a compound 3 or 4 wherein R_2 is 2,5-dioxo-1-imidazolidinyl having the 3-nitrogen atom unsubstituted can be alkylated at said atom to
5 obtain the corresponding N-alkyl derivatives, e.g. with methyl iodide in DMF in the presence of potassium carbonate. When R_1 and R_2 in the final compounds of formula (I) are, or contain, a reactive functional group, e.g. amino and hydroxy, these groups can be protected at the appropriate stage and and unmasked when convenient, according to
10 conventional methods well known in the art.

The chemistry in Scheme I and II proceeds through achiral or racemic intermediates, unless a substituent in R_1 or R_2 is chiral and enantiopure, or the aminoacid used in Scheme II for the preparation of compound 7 or 10 is chiral and enantiopure. When desired, pure enantiomers of the final products, in particular compounds of formula
15 (I') above, may be obtained from racemic compounds (I), or racemic intermediates thereof, by methods well known to one skilled in the art. For example, racemic carboxylates, including compounds (I) where X is HO- and intermediates 4, 9 where R is a hydrogen atom and n is zero, or intermediates 10 where R is a hydrogen atom, can be resolved by chiral salt formation, for example with enantiopure 1-phenylethylamine, brucine,
20 ephedrine, strychnine, morphine. Alternatively, final racemic compounds or racemic intermediates can be resolved by chiral chromatography. By another method, carboxylic esters, preferably methyl, ethyl and butyl esters, can be selectively hydrolysed to chiral acids by lipases and esterases. By a still different procedure, racemic carboxylic acids can be converted to diastereomeric esters with chiral alcohols such as borneol, menthol and the
25 like, which are then separated by crystallization or conventional chromatography. Alternatively, when in the acrylates of formula 2, 3, 8 in Schemes I and II R is the residue of a chiral alcohol such as borneol, menthol and the like, diastereomeric compounds 4, 9 10 are obtained, and variable degrees of diastereoselectivity can be achieved.

The present invention further provides a compound of formula (I) as defined above
30 for use in a method of treatment or prophylaxis of a disease mediated in a mammal by a matrix metalloproteinase, especially the gelatinases (MMP-2 and MMP-2), the membrane-

type MMP involved in gelatinase activation (MMP-14), the stromelysins (MMP-3 and MMP-10), collagenase-3 (MMP-13) and neutrophyl collagenase (MMP-8). Diseases for which the compounds of the present invention are particularly intended are: tumor growth and metastasis; rheumatoid arthritis and osteoarthritis; ophthalmic diseases, in particular
5 age-related macular degeneration and diabetic retinopathy; cardiovascular diseases, in particular restenosis; periodontal diseases; multiple sclerosis; Alzheimer disease.

The compounds of the present invention are typically administered in the form of pharmaceutical compositions, which can be prepared by combining the compounds of formula (I) or (I') or salts or hydrates or solvates thereof with a solid or liquid
10 pharmaceutically acceptable carrier, and, optionally, with pharmaceutically acceptable adjuvants and excipients, employing standard and conventional techniques. In particular, said compounds can be administered:

a) orally, for example, as tablets, coated tablets, capsules, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, or
15 syrups or elixirs. Compositions intended for oral use may be prepared according to any method known in the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active
20 ingredient in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, maize starch, or alginic acid; binding agents, for example starch, gelatin or acacia, and lubricating agents, for example
25 magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and adsorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed. Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed
30 with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is present as such, or mixed with water

or an oil medium, for example, peanut oil, liquid paraffin, or olive oil. Aqueous suspensions contain the active materials in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example, sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinylpyrrolidone gum tragacanth and gum acacia; dispersing or wetting agents may be naturally-occurring phosphatides, for example lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyoxyethylene sorbitan monooleate. The said aqueous suspensions may also contain one or more preservatives, for example, ethyl or n-propyl p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, or one or more sweetening agents, such as sucrose or saccharin. Oily suspension may be formulated by suspending the active ingredient in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents, such as those set forth above, and flavoring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an antioxidant such as ascorbic acid. Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, a suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present. The pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example olive oil or arachis oils, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally-occurring gums, for example gum acacia or gum tragacanth, naturally-occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides, for example sorbitan

mono-oleate, and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsion may also contain sweetening and flavoring agents. Syrups and elixirs may be formulated with sweetening agents, for example glycerol, sorbitol or sucrose. Such formulations may also contain a demulcent, a
5 preservative and flavoring and coloring agents;

b) parenterally, either subcutaneously, or intravenously, or intramuscularly, or intrasternally, or by infusion techniques, in the form of sterile injectable aqueous or oleaginous suspensions. This suspension may be formulated according to the known art using those suitable dispersing of wetting agents and suspending agents which have been
10 mentioned above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, for example as a solution in 1,3-butane diol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this
15 purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition fatty acids such as oleic acid find use in the preparation of injectables;

c) topically, in the form of creams, ointments, jellies, plasters, collyriums, solutions or suspensions;

d) rectally, in the form of suppositories prepared by mixing the drug with a suitable
20 non-irritating excipient which is solid at ordinary temperature but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials are cocoa butter and polyethylene glycols.

The compounds of this invention are advantageously administered orally in solid and liquid dosage forms. Daily dosages can vary within wide limits and will be adjusted to the
25 individual requirements in each particular case. As a rule of thumb, a compound of formula (I) or (I') is advantageously administered at a dosage and frequency to achieve and maintain, for the whole period of therapy, a blood level in free drug (i.e., not bound to plasma proteins) ranging from about 1 to about 100 times its K_i against the most relevant MMP for the particular disease state, e.g. MMP-2 for cancer. Thus, the daily dose may be divided into
30 multiple doses for administration, e.g. two to four times per day, which is generally expedient for compounds of formula (I) or (I') wherein X is HO-NH-; or, in the case of


particularly favorable pharmacokinetics, which can be found in compounds of formula (I) or (I') wherein X is HO-, the drug can be given daily, or even at alternate days. Generally, an amount of the active compound effective in adult patients will be in the range of 1 to about 500 mg, but it is to be understood that such range may be exceeded depending on the particular compound being used, the severity of the disease being treated, and the requirement of the patient.

The compounds of the present invention and their preparation will be better understood in connection with the following examples, which are intended as an illustration of and not a limitation upon the scope of the invention.

10

EXAMPLE 1

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfanyl)methyl]-propanoic acid.



Step 1. 4'-Chloro[1,1'-biphenyl]-4-sulfonic acid.

15

To a stirred solution of 4-chlorobiphenyl (21 g) in 150 mL of chloroform, cooled at -5 °C, was added dropwise chlorosulfonic acid (10.2 mL, 1.36 eq.). The cooling bath was removed and the mixture was stirred for 1 hour. Evaporation in vacuo left the title compound as a syrup.

Step 2. 4'-Chloro[1,1'-biphenyl]-4-sulfonyl chloride.

20

The crude product from step 1 was dissolved in thionyl chloride (100 mL) and refluxed for 1 hour. The solvent was removed in vacuo and the resulting solid was stripped twice with toluene (150 mL).

Step 3. 4'-Chloro[1,1'-biphenyl]-4-thiol.

25

4'-Chloro[1,1'-biphenyl]-4-sulfonyl chloride (30 g) was added portion wise to a stirred solution of tin(II) chloride dihydrate (80 g) in conc. HCl (50 mL) and water (150 mL) at 90-95 °C. The resulting yellow slurry was stirred for 1 hour at the same temperature. After

cooling, the precipitate was collected by filtration, thoroughly washed with water and dried to afford 21 g of bis[4-(4'-chloro[1,1'-biphenyl])] disulfide, m.p. 164-166 °C. This material was dissolved in THF (800 mL) and conc. HCl (25 mL), and Zn powder (20 g) was added under stirring. After stirring for 30 minutes at ambient temperature, the Zn was removed and rinsed with THF. The clear solution was concentrated in vacuo and partitioned between chloroform (450 mL) and brine. The organic phase was dried over anhydrous sodium sulfate and evaporated to leave 19 g of the title compound, mp 154-157 °C.

Step 4. 2-[[4'-Chloro[1,1'-biphenyl]-4-yl)sulfanyl]methyl]acrylic acid.

The thiol from step 3 (22 g) and 2-(bromomethyl)acrylic acid (18 g) were dissolved in 100 mL of DMF and stirred at 30 °C for 2 days under a low stream of nitrogen. The reaction mixture was diluted with ethyl acetate (400 mL) and washed several times with brine until neutral. The organic phase was dried over anhydrous sodium sulfate and evaporated in vacuo to obtain a crystalline mass. After addition of diethyl ether (150 mL) and swirling, the solid was collected by filtration, washed twice with ether, and dried to afford 21 g of the title compound, mp 191-195 °C.

Step 5. 2-[[4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]methyl]acrylic acid.

The material from step 4 above (20 g) was dissolved in 100 mL of DMF and added dropwise to a stirred solution of Oxone[®] (105 g) in 300 mL of water at 45-50 °C. After stirring for 5 hours, water (1 L) was added. The solid was collected by filtration, washed several times with water, then twice with ethanol (100 mL) and dried to provide 21.5 g of the title compound, mp 178-181 °C.

Step 6. 3-[[4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfanyl)methyl]propanoic acid.

Under a low stream of nitrogen, thiophenol (20 mL) was added to a stirred solution of the material from step 5 above (20 mg) in 35 mL of DMF. After stirring at 35 °C for 2 days, the reaction mixture was diluted with ethyl acetate and treated dropwise with a solution of iodine (21 g) in 75 mL of THF. The resulting brown solution was washed until complete decolorization with a solution of sodium metabisulfite (10 g) in 300 mL of water. After washing with brine and drying over anhydrous sodium sulfate, the solvent

was removed and the residue was filtered through a short pad of silica gel (450 g), eluting with chloroform, to obtain 23 g of the title compound, mp 158-162 °C.

¹H NMR (400 MHz, CDCl₃): δ 3.06 (m, 2H), 3.33 (m, 1H), 3.60 (dd, *J* = 3.4 and 14.2 Hz, 1H), 3.68 (dd, *J* = 7.3 and 14.2 Hz, 1H), 7.22 (m, 5H), 7.50 (m, 4H), 7.68 (d, *J* = 8.4 Hz, 2H), 7.88 (d, *J* = 8.4 Hz, 2H) [COOH: not seen]. MS (EI): *m/z* 446, 252, 194, 152, 149, 110. Anal. Found C, 59.32; H, 4.44; S, 14.32.

EXAMPLE 2

Preparation of sodium 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfanyl)methyl]-propanoate.

Sodium hydroxide (1 M, 48 mL) was added to a solution in methanol (75 mL) of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-(*RS*)-[(phenylsulfanyl)-methyl]propanoic acid (20 g, obtained as described in Example 1). Charcoal was added and the mixture was filtered. Concentration of the filtrate to 1/3 of the original volume furnished a shiny precipitate, which was collected by filtration and sequentially washed with cold water, ethanol and diethyl ether. Drying to a constant weight provided 19 g of the title compound as white crystals.

EXAMPLE 3

Preparation of (2*S*)-3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfanyl)methyl]-propanoic acid and of (2*R*)-3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfanyl)-methyl]propanoic acid.

Racemic 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfanyl)methyl]-propanoic acid, prepared as described in Example 1, was eluted over Chiralpak AS (Chiral Technologies) with ethanol + 0.05% trifluoroacetic acid (V/V) at 0.4 mL/min, monitoring the absorbance at 270 nm. The eluates were collected and concentrated in vacuo to yield the title enantiomers.

ISOMER 1, (2*S*) configuration:


R_f = 12.0 min (0.46x25 cm Chiralpak AS, ethanol + 0.05% TFA); [α]_D²⁵ = +47.8° (*c* = 0.83%, EtOH). Absolute configuration established by X-Ray crystallography.

ISOMER 2, (2*R*) configuration:

R_f = 15.0 min (0.46x25 cm Chiralpak AS, ethanol + 0.05% TFA); [α]_D²⁵ = -43° (*c* = 0.83%, EtOH).

EXAMPLE 4

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(phenylsulfanyl)-methyl]propanamide.




5 TBTU (0,72 g) was added portionwise to a stirred solution of N-methylmorpholine (0.5 mL) and O-tert-butylhydroxylamine hydrochloride (0.43 g) and 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfanyl)methyl]-propanoic acid (0.5 g) in 20 mL of DMF and 20 mL of acetonitrile cooled at -5°C. After stirring at ambient temperature overnight, the resulting solution was diluted with ethyl acetate and sequentially washed
10 with 0.1 M citric acid and 0.1 M NaHCO₃. After drying over anhydrous sodium sulfate and removal of the solvent, the crude reaction mixture was treated with trifluoroacetic acid (10 mL) and kept aside for 6 hours. The solvent was removed and the residue was partitioned between ethyl acetate and 0.1 M NaHCO₃. After drying, the solvent was evaporated off and the residue was chromatographed on silica gel eluting with
15 chloroform/cyclohexane 2/3 to afford, after crystallization from ethyl acetate, 0.35 g of the title compound, mp

¹H NMR (400 MHz, DMSO-*d*₆): δ 2.70(m, 1H), 3.09 (d, *J*= 7.8 Hz, 2H), 3.61 (m, 2H), 7.20 (m, 5H), 7.58 (d, *J*= 8.3 Hz, 2H), 7.78 (d, *J*= 8.3 Hz, 2H), 7.90 (m, 4H), 8.95 (s, 1H), 10.81 (s, 1H).. MS (EI): *m/z* 461, 400.

EXAMPLE 5

20 Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanoic acid.



Step 1. 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfanyl]-2-[(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanoic acid.

To a stirred solution of α -bromomethylacrylic acid (1.65 g) and NaHCO_3 (1.7 g) in 25 mL of DMF was added 3,4,4-trimethyl-2,5-dioxo-imidazolidine (2.13 g) and the resulting suspension was heated at 75°C for 3 h. After disappearance of α -bromomethylacrylic acid, checked by tlc and HPLC, the solution was cooled at $0-5^\circ\text{C}$ and 4'-chloro[1,1'-biphenyl]-4-thiol (3.3 g) was added. After stirring overnight at room temperature, the solution was diluted with ethyl acetate and washed thoroughly with 0.1 M citric acid. After drying and removal of the solvent, the residue was chromatographed on silica gel eluting with chloroform/cyclohexane 1/1 to provide, after crystallization from ethyl ether, 2.4 g of the title compound, mp $148-150^\circ\text{C}$.

Step 2. 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanoic acid.

A solution of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfanyl]-2-[(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanoic acid (4.6 g) in 30 ml of DMF was added dropwise to a stirred solution of Oxone[®] (13.5 g) in 75 mL of DMF and 20 mL of water at 45°C . After stirring for 2 h, the solvent was removed and the residue was partitioned between ethyl acetate and brine. The residue was columned on a small pad of silica gel eluting with chloroform to furnish 3.9 g of the title compound as a white foam.

^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 1.22(s, 3H) 1.24 (s,3H), 2.74 (s, 3H), 3.02 (m, 1H), 3.5-3.7 (m, 4H), 7.57 (d, $J=8.3$ Hz, 2H), 7.79 (d, $J=8.3$ Hz, 2H), 7.94 (m, 4H), 12.85 (broad s, 1H). MS (EI): m/z 478, 433, 252, 211, 152, 56. Anal. Found C, 54.89; H, 5.15; N, 5.64; S, 6.53.

EXAMPLE 6

Preparation of (2*R*)-3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanoic acid and of (2*S*)-3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanoic acid.

Racemic 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanoic acid, prepared as described in Example 5, was separated by chiral chromatography under the conditions described in Example 3.

ISOMER 1, (2*R*) configuration:


$R_f = 9$ min (0.46x25 cm Chiralpak AS, ethanol + 0.05% TFA); $[\alpha]_D^{25} = -8.7^\circ$ ($c = 0.83\%$, EtOH). The absolute configuration established by protein X-Ray in complex with truncated stromelysin (MMP-3).

ISOMER 2, (2*S*) configuration:

$R_f = 12$ min (0.46x25 cm Chiralpak AS, ethanol + 0.05% TFA); $[\alpha]_D^{25} = +8.2^\circ$ ($c = 0.83\%$, EtOH).

EXAMPLE 7

Preparation of 3-([1,1'-biphenyl]-4-ylsulfonyl)-2-[(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanoic acid.




Following the general procedure outlined in Example 5 and making non-critical variations but replacing 4'-chloro[1,1'-biphenyl]-4-thiol with [1,1'-biphenyl]-4-thiol in step 2, the title compound was obtained as a whitish crystalline powder, mp > 200 °C (dec.).

^1H NMR (400 MHz, DMSO- d_6): δ 1.16 (s, 3H), 1.17 (s, 3H), 2.68 (s, 3H), 2.96 (m, 1H), 3.43 (dd, $J = 3.6$ and 14.6 Hz, 1H), 3.54 (m, 2H), 3.63 (dd, $J = 8.7$ and 14.6 Hz, 1H), 7.40 (m, 3H), 7.70 (m, 2H), 7.87 (m, 4H), 12.90 (broad s, 1H). MS (EI): m/z 445, 427, 227, 209. Anal. Found C, 57.39; H, 5.66; N, 6.98; S, 5.95.

EXAMPLE 8

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanamide.




Following the general procedure outlined in Example 4 and making non-critical variations but replacing 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfonyl)methyl]propanoic acid with 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanoic acid, the title compound was obtained as a white crystalline powder.

¹H NMR (400 MHz, DMSO-*d*₆): δ 1.19 (s, 3H), 1.22 (s, 3H), 2.72 (s, 3H), 2.90 (m, 1H), 3.38 (dd, *J* = 2.9 and 14.6 Hz, 1H), 3.46 (m, 2H), 3.60 (dd, *J* = 8.3 and 14.6 Hz, 1H), 7.57 (d, *J* = 8.3 Hz, 2H), 7.78 (d, *J* = 8.3 Hz, 2H), 7.93 (m, 4H), 8.82 (s, 1H), 10.80 (s, 1H). ESI(+)-MS: *m/z* 494, 4.61, 235, 209. Anal. Found C, 43.58; H, 4.12; N, 6.61; S, 5.10.

EXAMPLE 9

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(4,4-dimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanoic acid.




Following the general procedure outlined in Example 5 and making non-critical variations but starting with 5,5-dimethyl-2,4-imidazolidinedione in step 1, the title compound was obtained as a white crystalline powder, mp 242-245 °C.

^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 1.21(s, 3H), 1.22 (s, 3H), 3.04 (m, 1H), 3.52 (m, 1H), 3.57 (m, 1H), 3.68 (m, 1H), 7.58 (d, $J=8.5$ Hz, 2H), 7.79 (d, $J=8.5$ Hz, 2H), 7.94 (m, 4H), 8.29 (s, 1H), 12.82 (broad s, 1H). ESI(+)-MS: m/z 465, 235, 213, 213, 195. Anal. Found C, 52.30; H, 4.82; N, 5.76; S, 6.44.

5

EXAMPLE 10

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(4,4-dimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanamide.



10

Following the general procedure outlined in Example 7 and making non-critical variations but starting in step 1 with 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(4,4-dimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanoic acid (prepared as described in Example 8), the title compound was obtained as a white crystalline powder.


15

^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 1.17(s, 3H), 1.21 (s, 3H), 3.34 (dd, $J=3.0$ and 15.0 Hz, 1H), 3.44 (m, 2H), 3.62 (dd, $J=9.0$ and 15.0 Hz, 1H), 7.58 (d, $J=8.9$ Hz, 2H), 7.78 (d, $J=8.9$ Hz, 2H), 7.93 (m, 4H), 8.24 (s, 1H), 8.83 (s, 1H), 10.81 (s, 1H). ESI(+)-MS: m/z 480, 447, 195. Anal. Found C, 54.06; H, 4.87; N, 8.67; S, 6.96.

EXAMPLE 11

20

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2,5-dioxo-1-imidazolidinyl)-methyl]propanoic acid.




Following the general procedure outlined in Example 5 and making non-critical variations but starting with 2,4-imidazolidinedione in step 1, the title compound was obtained as a white crystalline powder, mp 188-190 °C.

- 5 ¹H NMR (400 MHz, DMSO-*d*₆): δ 3.02 (m, 1H), 3.5-3.7 (m, 4H), 3.82 (d, *J*= 17.7 Hz, 1H), 3.87 (d, *J*= 17.7 Hz, 1H), 7.59 (d, *J*= 8.5 Hz, 2H), 7.80 (d, *J*= 8.5 Hz, 2H), 7.94 (m, 4H), 8.07 (s, 1H), 12.80 (broad s, 1H). ESI(+)-MS: *m/z* 437, 419, 235, 167. Anal. Found C, 47.71; H, 3.81; N, 5.79; S, 7.18.

EXAMPLE 12

- 10 Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(2,5-dioxo-1-imidazolidinyl)methyl]propanamide.




- 15 Following the general procedure outlined in Example 8 and making non-critical variations but starting in step 1 with 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2,5-dioxo-1-imidazolidinyl)methyl]propanoic acid (prepared as described in Example 11), the title compound was obtained as a yellowish crystalline powder, mp 150-156 °C.

- ¹H NMR (400 MHz, DMSO-*d*₆): δ 2.83 (m, 1H), 3.37 and 3.60 (each m, 1H), 3.42 (m, 2H), 3.76 (d, *J*= 17.6 Hz, 1H), 3.81 (d, *J*= 17.6 Hz, 1H), 7.57 (d, *J*= 8.3 Hz, 2H), 7.78 (d, *J*= 8.3 Hz, 2H), 7.91 (m, 4H), 8.01 (s, 1H), 8.70 (broad s, 1H), 10.80 (broad s, 1H).
20 ESI(+)-MS: *m/z* 452, 419, 235. Anal. Found C, 44.35; H, 3.87; N, 7.59; S, 6.81.

EXAMPLE 13

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3-butyl-2,5-dioxo-1-imidazolidinyl)methyl]propanoic acid.



Step 1. Methyl 2-[(3-butyl-2,5-dioxo-1-imidazolidinyl)methyl]acrylate.

Methyl 2-(bromomethyl)acrylate (0.7 mL) and 1-butyl-2,4-imidazolidinedione (1 g) were dissolved in DMF (3 mL). Sodium hydrogen carbonate (693 mg) was added, and the mixture heated at 80 °C for 2 hours, cooled to ambient temperature and partitioned between ethyl acetate and water. The organic layer was washed with water and brine, dried over anhydrous sodium sulfate and evaporated in vacuo to leave the title compound (1.0 g) as a colourless oil.

Step 2. Methyl 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfanyl]-2-[(3-butyl-2,5-dioxo-1-imidazolidinyl)methyl]propanoate.

The product from step 1 above (500 mg) and 4'-chloro[1,1'-biphenyl]-4-thiol (520 mg) were dissolved in DMF (5 mL). Sodium hydrogen carbonate (252 mg) was added, and the mixture heated at 60 °C for 1 hour, cooled to ambient temperature and partitioned between ethyl acetate and water. The organic layer was washed with water and brine, dried over anhydrous sodium sulfate and evaporated in vacuo to leave the crude title compound, which was used as such in the following step.

Step 3. Methyl 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3-butyl-2,5-dioxo-1-imidazolidinyl)methyl]propanoate.

The crude compound from step 2 above was taken up in DMF (7 mL). Oxone® (2 g) was added, and the mixture stirred for 4 hours at ambient temperature. Work-up as in step 1 left a residue, which was purified by flash-chromatography over silica (ethyl acetate – hexane), to obtain the title compound (650 mg) as a colorless foam.


Step 4. 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3-butyl-2,5-dioxo-1-imidazolidinyl)-methyl]propanoic acid.

2N Hydrochloric acid (5 mL) was added to a solution of the compound from step 3 above (650 mg) in acetic acid (2 mL), and the mixture was stirred at 80 – 90 °C. After 2 hours, 37% HCl (1 mL) was added and stirring at 80 °C was continued for additional 5 hours. The reaction mixture was evaporated in vacuo, and the residue partitioned between ethyl acetate and water. The organic layer was dried over anhydrous sodium sulfate and evaporated. The solvent was removed in vacuo and the resulting solid was stripped twice with toluene, to yield the title compound (420 mg) as a whitish powder.

¹H NMR (400 MHz, DMSO-*d*₆): δ 0.84 (t, *J* = 7.3 Hz, 3H), 1.07 (m, 2H), 1.41 (m, 2H), 2.99 (m, 1H), 3.20 (t, *J* = 7.1 Hz, 2H), 3.4-3.7 (m, 4H), 3.88 (d, *J* = 7.6 Hz, 1H), 3.92 (d, *J* = 7.6 Hz, 1H), 7.58 (d, *J* = 8.8 Hz, 2H), 7.80 (d, *J* = 8.8 Hz, 2H), 7.93 (m, 4H), 12.95 (broad s, 1H). ESI(+)-MS: *m/z* 515, 493, 475, 223. Anal. Found C, 57.39; H, 5.66; N, 6.98; S, 5.95.

EXAMPLE 14

Preparation of 3-[[4-(Benzoylamino)phenyl]sulfonyl]-2-[(2-thienylsulfanyl)methyl]propanoic acid.



Step 1. Bis(4-(benzoylamino)phenyl) disulfide.

Benzoyl chloride (8 mL) was added to a solution of bis(4-aminophenyl) disulfide (4 g) and TEA (10 mL) in 100 mL of dichloromethane, and the mixture was stirred for 2 hours at ambient temperature. The precipitate was collected by filtration and sequentially washed with water, diluted aqueous HCl and n-hexane to obtain the title compound as a white solid.

Step 2. N-(4-Sulfanylphenyl)benzamide.

Zn powder (0.5 g) and 2 N HCl (30 mL) was added in three portions under vigorous stirring to a suspension of the disulfide from step 1 in THF (100 mL). After 4 hour at ambient temperature, the mixture was filtered and the filtrate was concentrated in vacuo. The residue was taken up in ethyl acetate and sequentially washed with water and aqueous sodium hydrogen carbonate. Drying over anhydrous sodium sulfate and evaporation afforded 3.66 g of the title compound as a yellow solid.

Step 3. 2-[[[4-(Benzoylamino)phenyl]sulfanyl]methyl]acrylic acid.

A mixture of N-(4-sulfanylphenyl)benzamide (1.5 g), 2-(bromomethyl)acrylic acid (1 g) and TEA (1.75 mL) was stirred for 4 hours at ambient temperature. Washing with diluted aqueous HCl, drying over anhydrous sodium sulfate and evaporation in vacuo left a residue, which was passed through a short pad of silica gel (dichlorometane – methanol as the eluants) to obtain 1.68 g of the title compound as a colorless oil.

Step 4. 2-[[[4-(Benzoylamino)phenyl]sulfonyl]methyl]acrylic acid.

A mixture of Oxone[®] (10 g) and 2-[[[4-(benzoylamino)phenyl]sulfanyl]methyl]acrylic acid (1.68 g) in DMF (20 mL) was heated at 50 °C for 4 hours under vigorous stirring, then partitioned between ethyl acetate and 4% aqueous HCl. The organic phase was washed twice with water, dried over anhydrous sodium sulfate and evaporated to obtain the crude title compound as a yellowish oil.


Step 5. 3-[[4-(Benzoylamino)phenyl]sulfonyl]-2-[(2-thienylsulfanyl)methyl]propanoic acid.

2-Thiophenethiol (250 mg) was added to a solution of the crude acrylic acid from step 4 above (300 mg) in 5 mL of DMF. After 1 hour at ambient temperature, the mixture was partitioned between ethyl acetate and 4% aqueous HCl. The organic phase was washed twice with water, dried over anhydrous sodium sulfate and evaporated. The residue was chromatographed over silica gel with chloroform/methanol, and the eluate concentrated in vacuo to obtain 335 mg of the the title compound as a yellowish powder.

¹H NMR (400 MHz, DMSO-*d*₆): δ 2.70 (m, 1H), 2.98 (d, *J* = 7.1 Hz, 2H), 3.57 (m, 2H), 7.00 (m, 2H), 7.5-8.00 (m, 10H), 10.66 (s, 1H), 12.80 (broad s, 1H). ESI-MS: *m/z* 461, 260, 200, 182, 115, 105. Anal. Found C, 53.95; H, 4.65; N, 3.97; S, 18.55.

EXAMPLE 15

Preparation of 3-[[4-(Benzoylamino)phenyl]sulfonyl]-2-[(phenylsulfanyl)methyl]propanoic acid.



5


Following the general procedure outlined in Example 14 and making non-critical variations but starting with thiophenol in step 5, the title compound was obtained as a white powder.

¹H NMR (400 MHz, DMSO-*d*₆): δ 2.70 (m, 1H), 3.15 (m, 2H), 3.57 (dd, *J* = 4.4 and 14.4 Hz, 1H), 3.63 (dd, *J* = 8.1 and 14.4 Hz, 1H), 7.2-7.6 (m, 8H), 7.8-8.1 (m, 6H), 10.66 (s, 1H), 12.80 (broad s, 1H). ESI-MS: *m/z* 455, 331, 260, 194, 123, 110, 105, 77. Anal. Found C, 60.03; H, 5.29; N, 3.28; S, 11.68.

EXAMPLE 16

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[4-(hydroxyphenyl)sulfanyl]methyl]propanoic acid.

15



A solution of 2-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]methylacrylic acid (3.4 g) and 4-hydroxythiophenol (3 g) in 20 mL of DMF was heated for 5 h at 60°C. After cooling, the residue was chromatographed on silica gel eluting with CHCl₃/cyclohexane

20


1/2. The fractions containing the product were pooled and the solvent removed. The residue was crystallized from CHCl_3 /cyclohexane 1/3 affording 2.7 g of the title compound as a white powder.

^1H NMR (400 MHz, , DMSO- d_6): δ 2.58 (m, 1H), 2.95 (m, 2H), 3.64 (m, 2H), 6.63 (d, J = 8.5 Hz, 2H), 7.06 (d, J = 8.5 Hz, 2H), 7.58 (d, J = 8.5 Hz, 2H), 7.79 (d, J = 8.5 Hz, 2H), 7.86 (d, J = 8.5 Hz, 2H), 7.92 (d, J = 8.5 Hz, 2H), 9.60 (broad s, 1H), 12.80 (br s, 1H).

MS (EI): m/z 462, 251, 210, 152, 126. Anal. Found C, 56.95; H, 4.25; S, 13.68.

EXAMPLE 17

Preparation of 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-(phenoxy)methylpropanoic acid, sodium salt.



To a stirred solution of α -bromomethylacrylic acid (4.95 g) and NaHCO_3 (4 g) in 25 mL of DMF was added phenol (4.5 g) and the resulting suspension was heated at 75°C for 5 h. After disappearance of α -bromomethylacrylic acid, checked by tlc and HPLC, the solution was cooled at -15°C and 4'-chloro[1,1'-biphenyl]-4-thiol (9.2 g) was added. After stirring 1 h at room temperature, the solution was diluted with ethyl acetate and washed thoroughly with 0.1 M citric acid. After drying and removal of the solvent, the residue was chromatographed on silica gel eluting with CHCl_3 /cyclohexane 1/2 to provide 4.9 g of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfanyl]-2-phenoxyethylpropanoic acid as a white foam. A solution of this intermediate (4.6 g) in 50 mL of ethanol was added dropwise to a stirred solution of Oxone[®] (15 g) in 100 mL of DMF at 45°C . After stirring for 4 h, the solvent was evaporated off and the residue was partitioned between chloroform and water. After drying and removal of the solvent, the crude residue was chromatographed on silica gel eluting with chloroform to provide 3.9 g of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-


2-phenoxyethyl]propanoic acid as white foam. To a solution of this intermediate (2.15 g) in 100 mL of EtOH was added 1 M NaOH (6 mL). The resulting solution was charcoalized then filtered. Concentration in vacuum afforded 1.9 g of the title compound as shiny crystals.

5 ^1H NMR (400 MHz, DMSO- d_6): δ 2.55 (m, 1H), 3.44 (dd, J = 8.3 and 14.6 Hz, 1H), 3.97 (dd, J = 3.7 and 14.6 Hz, 1H), 4.01 (dd, J = 4.4 and 8.8 Hz, 1H), 4.20 (dd, J = 4.4 and 8.8 Hz, 1H), 6.75 (d, J = 8.8 Hz, 2H), 6.83 (t, J = 7.3 Hz, 1H), 7.18 (dd, J = 7.3 and 8.8 Hz, 2H), 7.56 (d, J = 8.8 Hz, 2H), 7.72 (d, J = 8.8 Hz, 2H), 7.85 (d, J = 8.5 Hz, 2H), 7.91 (d, J = 8.5 Hz, 2H). ESI(-)MS: m/z 429, 291, 251. Anal. Found C, 55.54; H, 4.52; S, 6.74.

10

EXAMPLE 18

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3-butyl-2,4-dioxo-1-imidazolidinyl)methyl]propanoic acid.



15

To stirred solution of glycine ethyl ester hydrochloride (6. g) and TEA (16 mL) in 80 mL of DMF was added α -bromomethylacrylic acid (5 g) at 0°C. After stirring for 2 h at rt, the suspension was filtered and the filtrate was evaporated to dryness. The residue dissolved in 50 mL of dioxane was treated with *n*-butyl isocyanide (4 mL) and heated for 5 h at 65°C. The solvent was removed and the crude residue was columned over a small pad of silica gel eluting with dichloromethane to afford 5.5 g of 2-[(3-butyl-2,4-dioxo-1-imidazolidinyl)-methyl]acrylic acid as an oil. This intermediate (5.5 g) was dissolved in DMF (50 mL) and treated at 0 °C with 4'-chloro[1,1'-biphenyl]-4-thiol (5.5 g) and NaHCO₃ (3.9 g). After stirring 1 h at rt, the reaction mixture was concentrated in vacuo, taken up in ethyl acetate, and sequentially washed with 0.5 M HCl and brine. After drying over anhydrous sodium sulfate, the solvent was removed and the residue was

20


25

chromatographed on silica eluting with ethyl acetate/cyclohexane 1/1 to provide 4.6 g of 3-
[(4'-chloro[1,1'-biphenyl]-4-yl)sulfanyl]-2-[(3-butyl-2,4-dioxo-1-
imidazolidinyl)methyl]propanoic acid. A solution of this intermediate (4.5 g) and Oxone[®]
(13 g) in 100 ml of DMF was heated for 1 h at 50°. The reaction mixture was concentrated
5 in vacuo and the residue was partitioned between ethyl acetate and water. After drying
over sodium sulfate, the solvent was removed in vacuo and the residue was columned over
a small pad of silica gel eluting with CHCl₃ to give the title compound (4.2 g) as a white
powder.

¹H NMR (400 MHz, DMSO-*d*₆): δ 0.83 (t, *J* = 7.3 Hz, 3H), 1.20 (m, 2H), 1.40 (m,
10 2H), 3.00 (m, 1H), 3.30 (m, 2H), 3.52 (d, *J* = 7.6 Hz, 2H), 3.53 (m, 1H), 3.71 (dd, *J* = 8.3
and 14.9 Hz, 1H), 3.79 and 3.93 (two d, *J* = 17.3 Hz, 2H), 7.58 (d, *J* = 8.5 Hz, 2H), 7.79 (d,
J = 8.5 Hz, 2H), 7.94 (s, 4H), 12.90 (broad s, 1H). ESI(+)-MS: *m/z* 493, 475, 319, 235,
223. Anal. Found C, 55.38; H, 5.19; N, 5.33; S, 6.78.

EXAMPLE 19

15 Preparation of 3-[(4'-cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfanyl)methyl]-
propanoic acid.



20 Step 1. 4-Cyano-4'-[[[(dimethylamino)carbothioyl]oxy]-1,1'-biphenyl].

Sodium hydride (80% in oil; 4.6 g) was added portionwise to a stirred solution of
4-hydroxy[1,1'-biphenyl]-4'-carbonitrile (26.8 g) in DMF (150 mL) at -10°C. After
stirring for 20 min, a solution of dimethylthiocarbamoyl chloride (23 g) in DMF (50 mL)
was added dropwise. The resulting mixture was maintained for additional 15 min at -
25 10°C, then heated at 80 °C for 1 h. After cooling, ice water was added. The resulting

precipitate was filtered, thoroughly washed with water, then with acetone, to furnish 37 g of the title compound as a white powder, mp 243-247 °C.

Step 2.. 4-Cyano-4'-[[dimethylamino)carbonyl]sulfanyl]-1,1-biphenyl.

The compound from step 1 above (33 g) was heated at 240 °C under a nitrogen
5 blanket. After melting, the compound was heated for additional 45 min, then cooled to room temperature and dissolved in acetone. Charcoal was added, filtered off, and the solvent removed in vacuo. The residue was crystallized from a small volume of boiling ethanol to obtain 30 g of the title compound, mp 268-270 °C.

Step 3. 4'-Cyano[1,1'-biphenyl]-4-thiol.

10 The compound from step 2 above (31 g) in methanol (1 L) was treated with 1N NaOH (120 mL) and the mixture was refluxed for 5 h. After cooling, 1 N HCl (150 mL) was added, and the mixture was concentrated in vacuo to remove the organic solvent. The crystalline precipitate was filtered, washed with water and then with ethanol, to provide 20 g of the title compound as white crystals, mp 192-195 °C.


15 Steps 4-6. 3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfanyl)methyl]propanoic acid.

By following the procedure described in Example 1, steps 4-6, and making non-critical variations, but replacing 4'-chloro[1,1'-biphenyl]-4-thiol with 4'-cyano[1,1'-biphenyl]-4-thiol, the title compound was obtained as a white powder.

20 ¹H NMR (400 MHz, DMSO-*d*₆): δ 3.09 (m, 2H), 3.36 (m, 1H), 3.61 (dd, *J*= 3.8 and 14.5 Hz, 1H), 3.70 (dd, *J*= 7.7 and 14.5 Hz, 1H), 7.20 (m, 5H), 7.6-7.7 (m, 6H), 7.94 (d, *J*= 8.5 Hz, 2H). Anal. Found C, 62.83; H, 4.41; N, 3.13; S, 14.29.

EXAMPLE 20

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2-pyridinylsulfanyl)methyl]-
25 propanoic acid.




Following the general procedure outlined in Example 16 and making non-critical variations but replacing 4-hydroxy-thiophenol with 2-mercaptopyridine, the title compound was obtained as a white crystalline powder.

¹H NMR (300 MHz, DMSO-*d*₆): δ 2.99 (m, 1H), 3.37 (d, *J* = 7.5 Hz, 1H), 3.66 (dd, *J* = 4.1 and 14.6 Hz, 1H), 3.70 (dd, *J* = 8.3 and 14.6 Hz, 1H), 7.01 (ddd, *J* = 1.0, 4.9 and 7.4 Hz, 1H), 7.18 (ddd, *J* = 0.9, 1.0 and 8.1 Hz, 1H), 7.50 (ddd, *J* = 1.9, 7.4 and 8.1 Hz, 1H), 7.56 (d, *J* = 8.6 Hz, 2H), 7.76 (d, *J* = 8.6 Hz, 2H), 7.88 (m, 4H), 8.30 (ddd, *J* = 0.9, 1.9 and 4.9 Hz, 1H), 12.85 (broad s, 1H). ESI(+)-MS: *m/z* 448, 319, 235, 178. Anal. Found C, 56.26; H, 4.09; N, 3.17; S, 14.30.

EXAMPLE 21

Preparation of 3-[[4-[(4-chlorobenzoyl)amino]phenyl]sulfonyl]-2-[(phenylsulfanyl)methyl]propanoic acid.




Following the general procedure outlined in Example 14 and making non-critical variations but starting with 4-chlorobenzoyl chloride in step 1, and replacing 2-thiophenethiol with thiophenol in step 5, the title compound was obtained as a white powder.

^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 2.67 (m, 1H), 3.15 (m, 2H), 3.62 (m, 2H), 7.22 (m, 5H), 7.64 (d, $J = 8.5$ Hz, 2H), 7.77 (d, $J = 8.0$ Hz, 2H), 8.01 (m, 4H), 10.70 (s, 1H), 12.80 (broad s, 1H). ESI(-)MS: m/z 510 $[\text{M} + \text{Na} - 2\text{H}]^+$, 334. Anal. Found C, 55.66; H, 4.10; N, 2.91; S, 12.82.

5

EXAMPLE 22

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2-pyrimidinylsulfanyl)methyl]-propanoic acid.



10 Step 1. 3-Bromo-2-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]methyl]propanoic acid.

2-[[[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]methyl]acrylic acid (4 g), prepared as described in Example 1, step 5, was suspended in 33% HBr in acetic acid (40 mL). After stirring for 4 h at ambient temperature, the reaction mixture was concentrated in vacuo, taken up in toluene and evaporated to dryness (twice). The residue was triturated with ethyl
15 ether, whereby the title product separated as a white crystalline powder, which was collected by filtration (3.85 g). A second crop (0.3 g) was recovered from the mother liquors by repeating the process after evaporation to dryness.

Step 2. 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2-pyrimidinylsulfanyl)methyl]-propanoic acid.


20 2-Mercaptopyrimidine (3.65 g) was added portionwise to a suspension of sodium hydride (60% dispersion in mineral oil; 1.21 g) in anhydrous DMF (80 mL) under an argon blanket, and the mixture stirred for 20 min at room temperature to obtain a clear solution, after which time the bromomethylpropanoic acid from step 1 above (4.54 g) was added. After 4 h at room temperature, 0.2 N HCl (600 mL) was added, and the mixture was
25 extracted twice with ethyl acetate. The organic extract was washed with brine, dried over sodium sulfate, evaporated, taken up in toluene and evaporated again to dryness.

Trituration of the residue with ethyl ether afforded a powder (2.99 g), which was purified by stirring for 1 h in a mixture of ether/ethyl acetate 9/1 and filtration, thereby obtaining the title product (2.7 g) as a white powder, mp 166-168 °C.

¹H NMR (300 MHz, DMSO-*d*₆): δ 3.06 (m, 1H), 3.28 (partially overlapped by water, 2H), 3.62 (dd, *J*= 4.0 and 14.6 Hz, 1H), 3.72 (dd, *J*= 8.2 and 14.6 Hz, 1H), 7.10 (t, *J*= 4.9 Hz, 1H), 7.57 (d, *J*= 8.5 Hz, 2H), 7.75 (d, *J*= 8.5 Hz, 2H), 7.90 (m, 4H), 8.50 (d, *J*= 4.9 Hz, 2H), 13.00 (broad s, 1H). ESI(+)MS: *m/z* 449, 319, 235, 179, 113. Anal. Found C, 53.65; H, 3.92; N, 6.21; S, 14.30.

EXAMPLE 23

10 Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[(5-amino-1,3,4-thiadiazol-2-yl)sulfanyl]methyl]propanoic acid.




Following the general procedure outlined in Example 22 and making non-critical variations but replacing 2-mercaptopyrimidine with 5-amino-2-mercapto-1,3,4-thiadiazole, the title compound was obtained as a yellowish powder, mp 184-186 °C.

¹H NMR (400 MHz, DMSO-*d*₆): δ 2.98 (m, 1H), 3.30 (m, 2H), 3.68 (m, 2H), 7.33 (s, 2H), 7.57 (d, *J*= 8.5 Hz, 2H), 7.78 (d, *J*= 8.5 Hz, 2H), 7.92 (m, 4H), 13.00 (broad s, 1H). ESI(+)MS: *m/z* 470, 337, 319, 235. Anal. Found C, 41.38; H, 3.42; N, 10.83; S, 22.89.

EXAMPLE 24

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2-methyl-2H-1,2,3-triazol-4-yl)sulfanyl]methyl]propanoic acid.




Following the general procedure outlined in Example 22 and making non-critical variations but replacing 2-mercaptopyrimidine with 4-mercapto-2-methyl-2H-1,2,3-triazole, the title compound was obtained as a white powder, mp 166-168 °C.

5 ¹H NMR (400 MHz, DMSO-*d*₆): δ 3.02 (m, 1H), 3.34 (m, 2H), 3.48 (m, 3H), 3.69 (m, 2H), 7.58 (d, *J* = 8.5 Hz, 2H), 7.79 (d, *J* = 8.5 Hz, 2H), 7.91 (m, 4H), 8.49 (s, 1H), 12.98 (broad s, 1H). ESI(+)-MS: *m/z* 452, 337, 319, 235. Anal. Found C, 50.18; H, 4.06; N, 9.25; S, 13.45.

EXAMPLE 25

10 Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[2-methyl-1H-imidazol-2-yl]sulfanyl]propanoic acid.




15 Following the general procedure outlined in Example 22 and making non-critical variations but replacing 2-mercaptopyrimidine with 2-mercapto-1-methyl-1H-imidazole, the title compound was obtained as a white powder.

¹H NMR (400 MHz, DMSO-*d*₆): δ 2.90 (m, 1H), 3.20 (m, 2H), 3.63 (m, 2H), 6.88 (d, *J* = 2.1 Hz, 1H), 7.16 (d, *J* = 2.1 Hz, 1H), 7.56 (d, *J* = 8.5 Hz, 2H), 7.78 (d, *J* = 8.5 Hz, 2H), 7.91 (m, 4H). ESI(+)-MS: *m/z* 451, 337, 235, 199, 181.

20

EXAMPLE 26

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[2,6-dimethyl-4-pyrimidinyl]sulfanyl]propanoic acid.




Following the general procedure outlined in Example 22 and making non-critical variations but replacing 2-mercaptopyrimidine with 2,6-dimethyl-4-mercaptopyrimidine, the title compound was obtained as a white powder, mp 170-172 °C.

ESI(+)-MS: m/z 477, 336, 251, 187, 152, 140. Anal. Found C, 55.05; H, 4.52; N, 4.64; S, 12.32.

EXAMPLE 27

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[4,6-dimethyl-2-pyrimidinyl)sulfanylmethyl]propanoic acid.




Following the general procedure outlined in Example 22 and making non-critical variations but replacing 2-mercaptopyrimidine with 4,6-dimethyl-2-mercaptopyrimidine, the title compound was obtained as a white powder, mp 170-172 °C.

¹H NMR (400 MHz, DMSO-*d*₆): δ 2.20 (s, 6H), 3.05 (m, 1H), 3.30 (m, 2H), 3.61 (dd, *J* = 3.4 and 14.5 Hz, 1H), 3.73 (d, *J* = 8.1 and 14.5 Hz, 1H), 6.81 (s, 1H), 7.58 (d, *J* = 8.5 Hz, 2H), 7.75 (d, *J* = 8.5 Hz, 2H), 7.88 (m, 4H), 12.90 (broad s, 1H). ESI(+)-MS: m/z 477, 319, 294, 235, 225, 141. Anal. Found C, 53.90; H, 4.52; N, 5.70; S, 12.48.

EXAMPLE 28

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(1,3-thiazol-2-ylsulfanylmethyl)propanoic acid.




Following the general procedure outlined in Example 22 and making non-critical variations but replacing 2-mercaptopyrimidine with 2-mercaptothiazole, the title compound was obtained as a white powder, mp 152-154 °C.

¹H NMR (400 MHz, DMSO-*d*₆): δ 3.00 (m, 1H), 3.45 (m, 2H), 3.63 (dd, *J* = 4.7 and 14.5 Hz, 1H), 3.74 (d, *J* = 7.7 and 14.5 Hz, 1H), 7.57 (m, 3H), 7.65 (d, *J* = 3.4 Hz, 1H), 7.60 (d, *J* = 8.5 Hz, 2H), 7.77 (d, *J* = 8.5 Hz, 2H), 7.91 (m, 4H), 13.00 (broad s, 1H). ESI(+)-MS: *m/z* 454, 337, 319, 294, 235, 184. Anal. Found C, 50.07; H, 3.69; N, 3.15; S, 21.14.

EXAMPLE 29

Preparation of 3-[(4'-cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-(phenoxymethyl)propanoic acid, sodium salt.




Following the general procedure outlined in Example 17 and making non-critical variations but replacing 4'-chloro[1,1'-biphenyl]-4-thiol with 4'-cyano[1,1'-biphenyl]-4-thiol, the title compound was obtained as a white powder.

Anal. Found C, 59.59; H, 4.45; N, 2.88; S, 6.85.

EXAMPLE 30

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(4-acetylaminophenoxy)methyl]propanoic acid.




Following the general procedure outlined in Example 17 and making non-critical variations but replacing phenol with 4-acetamidophenol, the title compound was obtained as a white powder.

Anal. Found C, 58.03; H, 4.61; N, 2.75; S, 6.84.

EXAMPLE 31


Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[4-(4-hydroxy-6-methyl-2-pyrimidinyl)sulfanyl]methyl]propanoic acid.



Following the general procedure outlined in Example 22 and making non-critical variations but replacing 2-mercaptopyrimidine with 4-hydroxy-6-methyl-2-mercaptopyrimidine, the title compound was obtained as a white powder, mp 165-168 °C.

EXAMPLE 32

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[1-methyl-1H-tetrazol-5-yl)sulfanyl]methyl]propanoic acid.




Following the general procedure outlined in Example 22 and making non-critical variations but replacing 2-mercaptopyrimidine with 1-methyl-1H-tetrazole-5-thiol, the title compound was obtained as a whitish powder, mp 98-100 °C.

¹H NMR (500 MHz, , DMSO-*d*₆): δ 3.08 (m, 1H), 3.54 (m, 2H), 3.70 (dd, *J*= 4.3 and 14.6 Hz, 1H), 3.76 (dd, *J*= 7.6 and 14.6 Hz, 1H), 3.85 (s, 3H), 7.59 (d, *J*= 8.5 Hz, 2H), 7.79 (d, *J*= 8.5 Hz, 2H), 7.92 (m, 4H), 13.00 (broad s, 1H). ESI(+)-MS: *m/z* 453, 337, 319, 294, 235. Anal. Found C, 47.64; H, 4.59; N, 13.18; S, 11.86.

EXAMPLE 33

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[2-(hydroxymethyl)phenoxy]-methyl]propanoic acid.




Following the general procedure outlined in Example 17 and making non-critical variations but replacing phenol with salicyl alcohol, the title compound was obtained as a crystalline powder.

Anal. Found C, 59.50; H, 4.62; S, 6.98.

EXAMPLE 34

Preparation of 3-[[4-[(4-chlorobenzoyl)amino]phenyl]sulfonyl]-2-(phenoxymethyl)-propanoic acid.




Following the general procedure outlined in Example 17 and making non-critical variations but replacing 4-chloro[1,1'-biphenyl]-4-thiol with 4-chloro-N-(4-sulfany)-benzamide, the title compound was obtained as a white powder.

Anal. Found C, 57.72; H, 4.28; N, 3.02; S, 6.62.

EXAMPLE 35

Preparation of 3-[[4-[(4-cyanobenzoyl)amino]phenyl]sulfonyl]-2-(phenoxy)methyl]-propanoic acid.




Following the general procedure outlined in Example 17 and making non-critical variations but replacing 4-chloro[1,1'-biphenyl]-4-thiol with 4-cyano-N-(4-sulfany)-benzamide, the title compound was obtained as a white powder.

Anal. Found C, 61.33; H, 4.26; N, 6.19; S, 7.15.

EXAMPLE 36

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(5-methyl-1,3,4-thiadiazol-2-yl)sulfanylmethyl]propanoic acid.

51




Following the general procedure outlined in Example 22 and making non-critical variations but replacing 2-mercaptopyrimidine with 5-methyl-2-mercapto-1,3,4-thiadiazole, the title compound was obtained as a withish powder, mp 147-150 °C.

5 Anal. Found C, 47.03; H, 3.74; N, 5.69; S, 18.73.

EXAMPLE 37

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(1H-benzimidazol-2-yl)sulfanylmethyl]-propanoic acid.



10

Following the general procedure outlined in Example 22 and making non-critical variations but replacing 2-mercaptopyrimidine with 2-mercapto-1H-benzimidazole, the title compound was obtained as a withish powder, mp 176-179 °C.

15 Anal. Found C, 55.43; H, 3.96; N, 5.57; S, 12.32.

EXAMPLE 38

Biochemical Activity Test

The inhibitory activity of the compounds of the present invention was evaluated in vitro against one or more of the MMP enzymes (gelatinases, collagenases, stromelysins) using the internal fluorescence quenching method.. The assay method is based on the hydrolysis of the MMP substrate, MCA-Pro-Leu-Gly-Leu-DPA-Ala-Arg-NH₂ at 37 °C (see Knight,

20

C.G. et al., FEBS Lett. 296:263-266, 1992). The enzymes cleave at the Gly-Leu bond removing the internally quenching DPA group. The release of the highly fluorescent peptide MCA-Pro-Leu was monitored with a LS50 Perkin Elmer fluorimeter equipped with a Peltier-thermostated cell holder using an excitation wavelength of 326 nm and an emission wavelength of 398 nm. Slits were set at 14/14. Proper determination of steady-state rates of substrate cleavage required a preincubation at 37 °C of 15 minutes to allow for complete equilibration of the enzyme-inhibitor complex. For the determination of the apparent K_i ($K_{i,app}$), the concentration of the inhibitor was varied at a constant and low concentration of the substrate (3 μ M). Rates of substrate hydrolysis, expressed as fluorescence arbitrary units per minute, were calculated by least-square analysis of the steady-state linear progression curves. The steady-state rates were fitted to the Michaelis equation describing competitive inhibition; in particular, $K_{i,app}$ were determined by fitting the data to the tight-binding equation of Morrison (Biochem. Biophys. Acta, 185:269-286, 1969) by non-linear methods. Since the Michaelis constants (K_m) for the MCA peptide substrate with the MMPs is quite high (70 μ M or greater; see Knight, C.G. et al., reference above) and by far exceeds the substrate concentration, the factor $(1 + S/K_m)$ can be approximated to unity without appreciable error, so that the determined $K_{i,app}$ will be essentially equal to K_i .

Human gelatinase-A (MMP-2) was obtained as pro-enzyme (72 kDa) and was activated with 1 mM 4-aminophenylmercuric acetate for 30 min at 37 °C immediately prior to use. The inhibitor was diluted into the assay from a stock solution in 100% DMSO, and controls substituted an equal volume of DMSO so that the final DMSO concentration from inhibitor and substrate dilution in all assays was 1%. The concentration of MMP-2 was 19.2 picoM. For compounds of formula (I) where X is -OH, the assay buffer was 50 mM Tris/HCl pH 6.5 containing 150 mM NaCl, 10 mM $CaCl_2$, 0.01 mM $ZnCl_2$ and 0.05% Brij 35 (Protocol A). Other conditions were as stated above.

Human fibroblast collagenase (MMP-1) was obtained as truncated recombinant enzyme encompassing residues 101-269 and did not require activation. Assay conditions were the same as above, but enzyme concentration was 200 picoM.

Inhibitory data (K_i 's against MMP-2 and MMP-1, all nanoM; Protocol A) for some representative compounds of formula (I) wherein X is -OH are displayed in Table 1.

TABLE 1

	Example No.	MMP-2 Ki (nanoM)	MMP-1 Ki (nanoM)
	1	8.3	> 2500
	3, Isomer 1	6.0	6600
5	3, Isomer 2	660	33190
	5	5.5	> 2000
	6, Isomer 1	3.7	3600
	6, Isomer 2	50	n.t.
	7	42	n.t.
10	9	3	n.t.
	11	4	n.t.
	13	12	5286
	14	8	n.t.
	15	7	81090
15	16	9	9697
	17	9.2	14740
	18	7.8	n.t.
	19	17	n.t.
	20	3.9	6291
20	21	1.4	50560
	22	8	8033
	23	25	n.t.
	24	24	n.t.
	25	16	n.t.
25	26	26.8	n.t.
	27	22.8	n.t.
	28	6.6	n.t.
	29	21.5	n.t.
	30	46	19580
30	32	26	n.t.
	33	15	n.t.

54

34	2.4	53980
35	4.8	67520
36	33.4	n.t.
37	18.2	n.t.

5

The compounds of formula (I) where X is –NHOH were tested against MMP-2 and MMP-1 under the conditions stated above, but the assay solution was buffered at pH 7.4 (Protocol B). Inhibitory data (K_i's, all picoM) for some representative compounds of formula (I) wherein X is –NHOH are displayed in Table 2.

10


TABLE 2

15

Example No.	MMP-2 K _i (picoM)	MMP-1 K _i (picoM)
4	159	149000
8	107	26800
10	50	12600
12	69	17100

CLAIMS

1. A compound which is a 3-arylsulfonyl-2-methyl propanoic acid derivative of formula (I):



wherein

X is HO-NH- or HO-;

10 R_1 is selected from phenyl, 4-chlorophenyl, 4-fluorophenyl, 4-cyanophenyl, benzamido (i.e., -NH-CO-Ph) and benzamido substituted on the terminal phenyl ring by C_1 - C_4 alkyl, fluoro, chloro, cyano or C_1 - C_4 alkoxy;

R_2 is selected from:

(a) -S-Ar or -S-CH₂-Ar, wherein Ar is a monocarbocyclic or bicarbocyclic aromatic moiety which is either unsubstituted or substituted with one or two substituents selected from C_1 - C_4 alkyl, phenyl, benzyl, C_1 - C_4 alkoxy, fluoro, chloro, bromo, nitro, cyano, hydroxy, amino, dimethylamino, acetamido, methylthio and acetyl;

15

(b) -O-Ar, wherein Ar is as defined above;

(c) -S-Het or -S-CH₂-Het, wherein Het is a heterocyclic ring selected from pyridine, pyrimidine, pyridazine, pyrazine, 1,2,5-triazine, imidazole, thiophene, furan, pyrrole, pyrazole, 1,3-thiazole, 1,3-oxazole, 1,2,3-triazole, 1,2,4-triazole, 1,3,4-thiadiazole, 1,3,4-oxadiazole, 1,2,3,4-tetrazole, quinoline, isoquinoline, indole, 1,3-benzoxazole, 1,3-benzothiazole, benzimidazole, [1,3]oxazolo[4,5-b]pyridine, [1,3]thiazolo[4,5-b]pyridine, [1,2,3,4]tetrazolo[1,5-b]pyridazine and purine, and wherein said Het group can be substituted with one to three substituents selected from C_1 - C_4 alkyl, phenyl, pyridyl,


20

25

benzyl, C₁₋₄ alkoxy, methylthio, fluoro, chloro, nitro, cyano, hydroxy, oxo, amino, methylamino, dimethylamino, 2-dimethylaminoethyl, acetamido and acetyl; and

- (d) 2,5-dioxo-1-imidazolidinyl or 2,4-dioxo-1-imidazolidinyl, either of which is optionally substituted at the carbon atom by one or two methyl, linear or branched C₂-C₄ alkyl, phenyl, benzyl or hydroxymethyl groups, and at the nitrogen atom with C₁-C₄ linear or branched alkyl;
or a pharmaceutically acceptable salt thereof.

2. A compound according to claim 1 which is an isomer having the configuration depicted in formula (I'):



- wherein X, R₁ and R₂ are as defined in claim 1.

3. A compound according to claim 1 which is selected from:

- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfonyl)methyl]propanoic acid;
(2S)-3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfonyl)methyl]propanoic acid;
3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(phenylsulfonyl)methyl]-propanamide;
3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[(4-hydroxyphenyl)sulfonyl]methyl]propanoic acid;
3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[[[(4-hydroxyphenyl)sulfonyl]-

- methyl]propanamide;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-(phenoxymethyl)propanoic acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-(phenoxymethyl)propanamide;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2-pyridinylsulfanyl)methyl]propanoic
5 acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(2-pyridinylsulfanyl)methyl]-
propanamide;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2-pyridinylmethyl)sulfanyl]methyl]-
propanoic acid;
- 10 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(2-pyridinylmethyl)sulfanyl]-
methyl]propanamide;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(5-methyl-1,3,4-thiadiazol-2-yl)sulfanyl]-
methyl]propanoic acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(5-methyl-1,3,4-thiadiazol-2-
15 yl)sulfanyl]methyl]propanamide;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(5-amino-1,3,4-thiadiazol-2-yl)-sulfanyl]-
methyl]propanoic acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(1-methyl-1H-imidazol-2-yl)sulfanyl]-
methyl]propanoic acid;
- 20 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2-pyrimidinylsulfanyl)methyl]propanoic
acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(1-methyl-1H-tetrazol-5-
yl)sulfanyl]methyl]-
propanoic acid;
- 25 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(1-benzyl-1H-tetrazol-5-
yl)sulfanyl]methyl]-
propanoic acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(1H-benzimidazol-2-yl)sulfanyl]methyl]-
propanoic acid;
- 30 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(1,3-thiazol-2-yl)sulfanyl]methyl]propanoic
acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[(2-methyl-2H-1,2,3-triazol-4-yl)sulfanyl]-methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[(5-phenyl-1,3,4-oxadiazol-2-yl)sulfanyl]-methyl]propanoic acid;

5 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[1-(3-pyridinyl)-1H-imidazol-2-yl)sulfanyl]-methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[(4-methyl-1,3-oxazol-2-yl)sulfanyl]methyl]-

10 propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[1,3]oxazolo[4,5-b]pyrimidin-2-yl-sulfanyl)-methyl]propanoic acid;

15 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[1,2,3,4]tetrazolo[1,5-b]pyridazin-6-yl-sulfanyl)methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[(2,6-dimethyl-4-pyrimidinyl)sulfanyl]-methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[(4,6-dimethyl-2-pyrimidinyl)sulfanyl]-methyl]propanoic acid;

20 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[(4-hydroxy-6-methyl-2-pyrimidinyl)-sulfanyl]methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[(4,6-dihydroxy-2-pyrimidinyl)sulfanyl]-methyl]propanoic acid;

25 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(9H-purin-6-ylsulfanyl)methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-(3-fluorophenoxymethyl)propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2-acetylaminophenoxy)methyl]propanoic acid;

30 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(4-acetylaminophenoxy)methyl]propanoic acid;

3-[(1,1'-Biphenyl)-4-ylsulfonyl]-2-[(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)methyl]-

propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)-

methyl]propanoic acid;

5 (2R)-3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanamide;

10 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(4,4-dimethyl-2,5-dioxo-1-imidazolidinyl)-methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(4,4-dimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanamide;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2,5-dioxo-1-imidazolidinyl)methyl]propanoic

15 acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(2,5-dioxo-1-imidazolidinyl)-methyl]propanamide;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3-methyl-2,5-dioxo-1-imidazolidinyl)-methyl]propanoic acid;

20 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3-ethyl-2,5-dioxo-1-imidazolidinyl)methyl]-propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3-butyl-2,5-dioxo-1-imidazolidinyl)methyl]-

25 propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(3-butyl-2,5-dioxo-1-imidazolidinyl)methyl]propanamide;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3-butyl-2,4-dioxo-1-imidazolidinyl)methyl]-

30 propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(3-butyl-2,4-dioxo-1-

- imidazolidinyl)methyl]propanamide;
3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(5-isopropyl-2,4-dioxo-1-imidazolidinyl)-
methyl]propanoic acid;
3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(5-hydroxymethyl-2,4-dioxo-1-
5 imidazolidinyl)methyl]propanoic acid;
3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(5-hydroxymethyl-3-methyl-2,4-dioxo-1-
imidazolidinyl)methyl]propanoic acid;
3-[(4'-Fluoro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfanyl)methyl]propanoic acid;
3-[(4'-Fluoro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(phenylsulfanyl)methyl]-
10 propanamide;
3-[(4'-Fluoro[1,1'-biphenyl]-4-yl)sulfonyl]-2-(phenoxymethyl)propanoic acid;
3-[(4'-Fluoro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-(phenoxymethyl)propanamide;
3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfanyl)methyl]propanoic acid;
3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(phenylsulfanyl)methyl]-
15 propanamide;
3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-(phenoxymethyl)propanoic acid;
3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-(phenoxymethyl)propanamide;
3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-(3-fluorophenoxymethyl)propanoic acid;
3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2-acetylamino)phenoxymethyl]propanoic
20 acid;
3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2-pyridinylsulfanyl)methyl]propanoic acid;
3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[2-(2-pyridinylmethyl)sulfanyl]methyl]-
propanoic acid;
3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3-butyl-2,5-dioxo-1-
25 imidazolidinyl)methyl]-
propanoic acid;
3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(3-butyl-2,5-dioxo-1-
imidazolidinyl)methyl]propanamide;
3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(5-hydroxymethyl-2,4-dioxo-1-
30 imidazolidinyl)methyl]propanoic acid;
3-[[4-(Benzoylamino)phenyl]sulfonyl]-2-[(phenylsulfanyl)methyl]propanoic acid;

- 3-[[4-(Benzoylamino)phenyl]sulfonyl]-N-hydroxy-2-[(phenylsulfanyl)methyl]-propanamide;
- 3-[[4-(Benzoylamino)phenyl]sulfonyl]-2-[[2-(pyridinylmethyl)sulfanyl)methyl]propanoic acid;
- 5 3-[[4-(Benzoylamino)phenyl]sulfonyl]-2-[(2-thienylsulfanyl)methyl]propanoic acid;
- 3-[[4-(Benzoylamino)phenyl]sulfonyl]-2-(phenoxymethyl)propanoic acid;
- 3-[[4-[(4-Methylbenzoyl)amino]phenyl]sulfonyl]-2-[(phenylsulfanyl)methyl]propanoic acid;
- 3-[[4-[(4-Methylbenzoyl)amino]phenyl]sulfonyl]-N-hydroxy-2-[(phenylsulfanyl)methyl]-propanamide;
- 10 3-[[4-[(4-Chlorobenzoyl)amino]phenyl]sulfonyl]-2-[(phenylsulfanyl)methyl]propanoic acid;
- 3-[[4-[(4-Chlorobenzoyl)amino]phenyl]sulfonyl]-N-hydroxy-2-[(phenylsulfanyl)methyl]-propanamide;
- 15 3-[[4-[(4-Chlorobenzoyl)amino]phenyl]sulfonyl]-2-(phenoxymethyl)propanoic acid;
- 3-[[4-[(4-Chlorobenzoyl)amino]phenyl]sulfonyl]-N-hydroxy-2-(phenoxymethyl)-propanamide;
- 3-[[4-[(4-Chlorobenzoyl)amino]phenyl]sulfonyl]-2-(3-fluorophenoxymethyl)propanoic acid;
- 20 3-[[4-[(4-Chlorobenzoyl)amino]phenyl]sulfonyl]-2-[[2-(pyridinylmethyl)sulfanyl)methyl]-propanoic acid;
- 3-[[4-[(4-Chlorobenzoyl)amino]phenyl]sulfonyl]-2-[[1-methyl-1H-tetrazol-5-yl)sulfanyl]-methyl]propanoic acid;
- 3-[[4-[(4-Chlorobenzoyl)amino]phenyl]sulfonyl]-2-[(5-hydroxymethyl-2,4-dioxo-1-imidazolidinyl)methyl]propanoic acid;
- 25 3-[[4-[(4-Cyanobenzoyl)amino]phenyl]sulfonyl]-2-[(phenylsulfanyl)methyl]propanoic acid;
- 3-[[4-[(4-Cyanobenzoyl)amino]phenyl]sulfonyl]-N-hydroxy-2-[(phenylsulfanyl)methyl]-propanamide;
- 30 3-[[4-[(4-Cyanobenzoyl)amino]phenyl]sulfonyl]-2-[[1-methyl-1H-tetrazol-5-yl)sulfanyl]-methyl]propanoic acid;

3-[[4-[(4-Cyanobenzoyl)amino]phenyl]sulfonyl]-2-(phenoxymethyl)propanoic acid;

3-[[4-[(4-Cyanobenzoyl)amino]phenyl]sulfonyl]-N-hydroxy-2-(phenoxymethyl)-
propanamide;

3-[[4-[(4-Cyanobenzoyl)amino]phenyl]sulfonyl]-2-(3-fluorophenoxymethyl)propanoic
5 acid;

3-[[4-[(4-Cyanobenzoyl)amino]phenyl]sulfonyl]-N-hydroxy-2-(3-fluorophenoxymethyl)-
propanamide;

3-[[4-[(4-Cyanobenzoyl)amino]phenyl]sulfonyl]-2-[(2-acetylamino)phenoxymethyl]-
propanoic acid;


10 3-[[4-(4-Propylbenzoylamino)phenyl]sulfonyl]-2-[[[(1-methyl-1H-tetrazol-5-yl)sulfanyl]-
methyl]propanoic acid; and

3-[[4-[(4-Methoxybenzoyl)amino]phenyl]sulfonyl]-2-(phenoxymethyl)propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[2-(hydroxymethyl)phenoxy]methyl]-
propanoic acid; and the pharmaceutically acceptable salts thereof.

15

4. A process for producing a compound as defined in claim 1, starting from a
compound of formula 4:



20 wherein R is H or the residue of a carboxylic acid ester, R₁ and R₂ are as defined in
claim 1 and n is 0 or 2, which process comprises:

(a) hydrolysing a said compound of formula 4 in which R is the residue
of a carboxylic and ester to give a compound of formula (I) in which X is HO-; or

(b) hydrolysing and oxidising, in either order, a said compound of
25 formula 4 in which n is 0 and R is the residue of a carboxylic acid ester, to give a
compound of formula (I) in which X is HO-; or


(c) activating a said compound of formula 4 wherein R is H and n is 2 to form an activated carboxy group, coupling the activated carboxy group with hydroxylamine or an O-protected derivative thereof and, if necessary, deprotecting the hydroxamic group to give a compound of formula (I) wherein X is -NHOH ; or

(d) submitting a said compound of formula 4 wherein R is H and n is zero to a sequence of reactions comprising oxidation at the sulphur atom, activation of the carboxy group, condensation of the activated carboxy group with hydroxylamine or an O-protected derivative thereof and, if necessary, deprotection of the hydroxamic group to form a compound of formula (I) wherein X is -NHOH , the oxidation step being conducted either before the activation step or after the condensation step; and/or

(e) if desired, converting a resulting compound of formula (I) into another compound of formula (I); and/or converting a free compound into a pharmaceutically acceptable salt thereof; and/or converting a salt into a free compound.


5. A process according to claim 4 wherein the compound of formula 4 is obtained by

(a) subjecting a compound of formula 2:




wherein R, R_1 and n are as defined in claim 4, to conjugate addition by treatment with a compound of formula $R_2\text{H}$ wherein R_2 is as defined in claim 1; or

(b) treating a compound of formula 3:



wherein R and R_2 are as defined in claim 4, with a thiol of formula:



to obtain a compound of formula 4 in which n is zero.

- 5 6. A pharmaceutical composition which comprises a pharmaceutically acceptable carrier or diluent and, as an active principle, a compound as defined in any one of claims 1 to 4.
7. A compound as defined in any one of claims 1 to 4, for use in a method of
10 treatment of the human or animal body by therapy.
8. A compound as claimed in claim 7 for use in the treatment or prophylaxis of a disease in a mammal mediated by a matrix metalloproteinase.
- 15 9. A compound as claimed in claim 8 wherein the matrix metalloproteinase is a gelatinase (MMP-2), the membrane-type MMP involved in gelatinase activation (MMP-14), a stromelysin (MMP-3 or MMP-10), collagenase (MMP-13) or neutrophil collagenase (MMP-8).
- 20 10. A compound as claimed in any one of claims 7 to 9 for use in the treatment or prophylaxis of tumour growth or metastasis, rheumatoid arthritis, osteoarthritis, ophthalmic disease, cardiovascular disease, periodontal disease, multiple sclerosis or Alzheimer's disease.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 00/06429

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07C317/46 C07C323/65 C07D213/70 C07D233/74 C07D233/84
 C07D235/28 C07D239/38 C07D249/12 C07D257/04 C07D277/36
 C07D285/12 C07D333/34 A61K31/12 A61K31/16 A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

BEILSTEIN Data, WPI Data, EPO-Internal, PAJ, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 98 13340 A (PHARMACIA & UPJOHN) 2 April 1998 (1998-04-02) cited in the application page 2 -page 5; claims 1,11-18; examples 13-16,18-21	1,6-10
Y	WO 98 05635 A (CHIROSCIENCE) 12 February 1998 (1998-02-12) page 2 -page 4; claims 1,18-22; examples 24,56,A-D	1,6-10

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- * & * document member of the same patent family

Date of the actual completion of the international search

6 December 2000

Date of mailing of the international search report

19/12/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

English, R

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 00/06429

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61P19/02 A61P9/00 A61P25/28

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

6 December 2000

Date of mailing of the international search report

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

English, R

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 00/06429

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9813340 A	02-04-1998	AU 4645997 A	17-04-1998
		BR 9712134 A	31-08-1999
		CN 1230177 A	29-09-1999
		CZ 9900923 A	14-07-1999
		EP 0929519 A	21-07-1999
		HU 0000145 A	28-07-2000
		NO 991494 A	26-05-1999
		PL 332509 A	13-09-1999
		US 5847153 A	08-12-1998
WO 9805635 A	12-02-1998	AU 3856497 A	25-02-1998
		BR 9711027 A	17-08-1999
		CN 1227540 A	01-09-1999
		CZ 9900368 A	14-07-1999
		EP 0968182 A	05-01-2000
		NO 990543 A	06-04-1999
		PL 331598 A	02-08-1999
		US 6118001 A	12-09-2000
		ZA 9707044 A	07-08-1998